



PATIENT & CAREGIVER EDUCATION

Tigecycline

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Tygacil

Brand Names: Canada

Tygacil

Warning

- A larger number of deaths happened in people taking this drug compared to people taking other drugs used to treat infections. The cause of this is not known. This drug is only for use when other drugs cannot be used.
- Some products have maltose in them, which may affect certain glucose tests. False results of glucose tests may cause problems when treating high or low blood sugar. This may lead to long-lasting effects or death. Talk with the doctor about which blood sugar tests are best to use.

What is this drug used for?

- It is used to treat bacterial infections.

Children:

- This drug is not approved for use in children. However, the doctor may decide the benefits of taking this drug outweigh the risks. If your child has been given this drug, ask the doctor for information about the benefits and risks. Talk with the doctor if you have questions about giving this drug to your child.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have a hole in the GI (gastrointestinal) tract.
- If you are breast-feeding. Do not breast-feed if you will be taking this drug for longer than 3 weeks. You will need to talk about the benefits and risks to you and the baby.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

For all patients taking this drug:

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Have your blood work and other lab tests checked as

you have been told by your doctor.

- Do not use longer than you have been told. A second infection may happen.
- This drug may make you sunburn more easily. Use care if you will be in the sun. Tell your doctor if you sunburn easily while taking this drug.
- This drug may change tooth color to yellow-gray brown if taken by children younger than 8 years old, or in the unborn baby if taken during some parts of pregnancy. If this change of tooth color happens, it will not go away. Other tooth problems have also happened. Bone growth may also be affected in these people taking this drug. If you have questions, talk with the doctor.
- Birth control pills and other hormone-based birth control may not work as well to prevent pregnancy. Use some other kind of birth control also like a condom when taking this drug.
- Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Rarely, some allergic reactions have been life-threatening.
- Signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Signs of too much acid in the blood (acidosis) like confusion; fast breathing; fast heartbeat; a heartbeat that does not feel normal; very bad stomach pain, upset stomach, or throwing up; feeling very sleepy; shortness of breath; or feeling very tired or weak.
- Feeling very tired or weak.
- Not able to pass urine or change in how much urine is

passed.

- A round gray, brown, black, pink, or tan spot on the skin. Other spots can also appear.
- Diarrhea is common with antibiotics. Rarely, a severe form called C diff–associated diarrhea (CDAD) may happen. Sometimes, this has led to a deadly bowel problem. CDAD may happen during or a few months after taking antibiotics. Call your doctor right away if you have stomach pain, cramps, or very loose, watery, or bloody stools. Check with your doctor before treating diarrhea.
- Severe and sometimes deadly pancreas problems (pancreatitis) have happened with this drug. Call your doctor right away if you have severe stomach pain, severe back pain, or severe upset stomach or throwing up.
- Raised pressure in the brain has happened with drugs like this one. Most of the time, this went back to normal after the drug was stopped. Sometimes, loss of eyesight happened and did not go away even after the drug was stopped. Call your doctor right away if you have a headache or eyesight problems like blurred eyesight, seeing double, or loss of eyesight.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Diarrhea, stomach pain, upset stomach, or throwing up.
- Headache.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1- 800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- It is given as an infusion into a vein over a period of time.

What do I do if I miss a dose?

- Call your doctor to find out what to do.

How do I store and/or throw out this drug?

- If you need to store this drug at home, talk with your doctor, nurse, or pharmacist about how to store it.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information

leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.

- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a

specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2025-05-01

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Tigecycline - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center

Last Updated

December 12, 2022

Learn about our [Health Information Policy](#).