



PATIENT & CAREGIVER EDUCATION

Typhoid Vaccine

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Typhim Vi; Vivotif

Brand Names: Canada

Typhim Vi; Vivotif

What is this drug used for?

- It is used to prevent typhoid fever.

What do I need to tell my doctor BEFORE I take this drug?

All products:

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have an infection or an illness with a fever.
- If you have a weak immune system.
- If you are taking any drugs to suppress your immune system. This may be certain doses of steroids like prednisone. There are many drugs that can suppress your immune system. Ask your doctor or pharmacist if you are not sure.
- If you have typhoid fever or you carry typhoid in your blood.

Capsules:

- If you have a stomach illness like diarrhea or throwing up.
- If you are taking any of these drugs: Antibiotic, proguanil, or a sulfa drug. Ask your doctor or pharmacist if you are not sure.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Vaccines may not fully protect all people who get them. If you have questions, talk with the doctor.
- Use care when eating food and drinking water in disease-affected areas. If you have questions, talk with your doctor.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

All products:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.

Injection:

- Severe dizziness or passing out can happen after or sometimes before getting a vaccine. Tell your doctor right away if you feel dizzy.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

Injection:

- Pain, redness, swelling, or other reaction where the injection was given.
- Feeling tired or weak.
- Muscle pain.
- Upset stomach.
- Headache.

Capsules:

- Stomach pain.
- Upset stomach.
- Headache.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

Report side effects to the FDA/CDC Vaccine Adverse Event Reporting System (VAERS) at <https://vaers.hhs.gov/reportevent.html> or by calling 1-800-822-7967.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

Capsules:

- For this drug to work the best, be sure to take all of your doses when you were told by your doctor.
- Swallow this drug as soon as you can after putting it in your mouth. Take about 1 hour before a meal with a cold or lukewarm (not hot) drink.
- Do not chew this drug.
- All doses of this vaccine must be taken at least 1 week before you will be exposed to the bacteria that causes typhoid fever (*S. typhi*). If you cannot finish this vaccine 1 week before exposure, talk with your doctor.

Injection:

- It is given as a shot into a muscle.

What do I do if I miss a dose?

- Call your doctor to find out what to do.

How do I store and/or throw out this drug?

Capsules:

- Store capsules in original container in a refrigerator.
- Do not freeze.

Injection:

- If you need to store this drug at home, talk with your doctor, nurse, or pharmacist about how to store it.

All products:

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Additional Information

Vaccine Information Statements (VIS) are made by the staff of the Centers for Disease Control and Prevention (CDC). Each VIS gives information to properly inform the adult receiving the vaccine or, in the case of a minor, the child's parent or legal representative about the risks and benefits of each vaccine.

Before a doctor vaccinates a child or an adult, the provider is required by the National Childhood Vaccine Injury Act to give a copy of the VIS. You can also get foreign language versions.

<https://www.cdc.gov/vaccines/hcp/current-vis/typhoid.html>

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2026-02-25

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Typhoid Vaccine - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center