



PATIENT & CAREGIVER EDUCATION

Vaccinia Immune Globulin (Intravenous)

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US
CNJ-016

Warning

- This drug may affect certain glucose tests. False results of glucose tests may lead to not treating low blood sugar or to the use of more insulin. This may lead to long-lasting or life-threatening effects.
- If you have high blood sugar (diabetes), talk with your doctor about which glucose tests are best to use.

What is this drug used for?

- It is used to treat certain reactions or infections caused by the vaccinia virus.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have a certain eye infection caused by the vaccinia virus.
- If you have IgA deficiency.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists,

and dentists.

- This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug.
- This drug is made from human plasma (part of the blood) and may have viruses that may cause disease. This drug is screened, tested, and treated to lower the chance that it carries an infection. Talk with the doctor.
- Talk with your doctor before getting any vaccines. Use of some vaccines with this drug may either raise the chance of an infection or make the vaccine not work as well.
- Kidney problems have happened with human immune globulin. Sometimes, these problems have been deadly. Kidney problems are more common in people using products that have sucrose. Most immune globulin products do not have sucrose. The chance of these problems may be raised if you have kidney problems, high blood sugar (diabetes), fluid loss (dehydration) or low blood volume, a blood infection, or proteins in the blood that are not normal. The chance may be raised if you are 65 or older, or if you take other drugs that may harm the kidneys. Talk with your doctor if you have questions about this information or about if your product has sucrose.

- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Feeling very tired or weak.
- Dark urine or yellow skin or eyes.
- Severe dizziness or passing out.
- Pain where the shot was given.
- Fever or chills.

- Joint pain.
- A burning, numbness, or tingling feeling that is not normal.
- Pale skin.
- Severe lung problems have happened with this drug. This type of lung problem can be life-threatening. Call your doctor right away if you have lung or breathing problems like trouble breathing, shortness of breath, or a cough that is new or worse.
- This drug may raise the chance of a severe brain problem called aseptic meningitis. Call your doctor right away if you have a headache, fever, chills, very upset stomach or throwing up, stiff neck, rash, bright lights bother your eyes, feeling sleepy, or feeling confused.
- There is a chance of blood clots with this drug. The chance is raised in older people and in people with thick blood, heart problems, or a history of blood clots. The chance may also be higher if you must be in a bed or chair for some time, if you take estrogen, or if you use certain catheters. However, blood clots can happen even if you do not have any of these health problems. Call your doctor right away if you have numbness or weakness on 1 side of your body; pain, redness, tenderness, warmth, or swelling in the arms or legs;

change in color of an arm or leg; chest pain or pressure; shortness of breath; fast heartbeat; or coughing up blood. Talk with your doctor.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Feeling dizzy, tired, or weak.
- Feeling hot or cold.
- Headache.
- Upset stomach or throwing up.
- Decreased appetite.
- Muscle spasm.
- Back pain.
- Shakiness.
- Sweating a lot.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1- 800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- It is given as an infusion into a vein over a period of time.

What do I do if I miss a dose?

- Call your doctor to find out what to do.

How do I store and/or throw out this drug?

- If you need to store this drug at home, talk with your doctor, nurse, or pharmacist about how to store it.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do

so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the

medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2025-08-26

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors.

All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Vaccinia Immune Globulin (Intravenous - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center