PATIENT & CAREGIVER EDUCATION

Amphetamine

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Adzenys XR-ODT; Dyanavel XR; Evekeo; Evekeo ODT [DSC]

Brand Names: Canada

Dyanavel XR

Warning

- This drug has a high risk of misuse. This can lead to alcohol or drug use disorder. Misuse or abuse of this drug can lead to overdose or death. The risk is higher with higher doses or when used in ways that are not approved like snorting or injecting it. Do not give this drug to anyone else. This drug may also be habit-forming if taken for a long time. Do not give for longer than you have been told by the doctor. Give only as you were told. Tell the doctor if your child has ever had alcohol or drug use disorder. Your child will be watched closely while taking this drug. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs as you have been told.

What is this drug used for?

- It is used to treat attention deficit problems with hyperactivity.
- It is used to treat narcolepsy.
- It is used for weight loss.
- It may be given to your child for other reasons. Talk with the doctor.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child or a family member has any of these health problems: Blood vessel disease, high blood pressure, heart structure problems or other heart problems, or Tourette's syndrome or tics.
- If your child has any of these health problems: Glaucoma, agitation, anxiety, or overactive thyroid.
- If your child has ever had a stroke.
- If your child is taking any of these drugs: Acetazolamide or sodium bicarbonate.
- If your child has taken certain drugs for depression or certain other health problems in the last 14 days. This includes isocarboxazid, phenelzine, or tranylcypromine. Very high blood pressure may happen.
- If your child is taking any of these drugs: Linezolid or methylene blue.

If your child is breast-feeding a baby:

- Be sure your child does not breast-feed a baby while taking this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or

OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

All products:

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child avoid tasks or actions that call for alertness or clear eyesight until you see how this drug affects your child. These are things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized vehicles.
- Have your child's blood work and heart function checked as you have been told by the doctor.
- This drug may cause high blood pressure.
- Check blood pressure and heart rate as the doctor has told you.
- This drug may affect certain lab tests. Tell all of your child's health care providers and lab workers that your child takes this drug.
- Alcohol may interact with this drug. Be sure your child does not drink alcohol.
- If your child has been taking this drug for a long time or at high doses, it may not work as well and your child may need higher doses to get the same effect. This is known as tolerance. Call the doctor if this drug stops working well. Do not give more than ordered.
- Long-term or regular use of this drug may lead to dependence. Stopping this drug all of a sudden may lead to signs of withdrawal. Talk to your child's doctor before you lower the dose or stop giving this drug. You will need to follow the doctor's instructions. Tell your child's doctor if your child has any bad effects.

- Talk with the doctor before giving OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen or naproxen, and some natural products or aids.
- This drug may raise the chance of seizures in some people, including people who have had seizures in the past. Talk to the doctor to see if your child has a greater chance of seizures while taking this drug.
- This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor.
- Different brands of this drug may be for use in different ages of children. Talk with the doctor before giving this drug to a child.

If your child is pregnant:

- Tell the doctor if your child is pregnant or becomes pregnant. You will need to talk about the benefits and risks of your child using this drug while pregnant.
- Taking this drug during pregnancy may lead to withdrawal in the newborn.

All extended-release products:

- The risk of weight loss and other side effects is higher in children younger than 6 years of age. This drug is not for use in children younger than 6 years of age.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or

swelling of the mouth, face, lips, tongue, or throat.

- Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Signs of a urinary tract infection (UTI) like blood in the urine, burning or pain when passing urine, feeling the need to pass urine often or right away, fever, lower stomach pain, or pelvic pain.
- Shakiness.
- Change in eyesight.
- Seizures.
- Trouble controlling body movements.
- Trouble controlling sounds, like humming, throat clearing, yelling, or making loud noises.
- Change in color of hands, feet, or other areas. Skin may turn pale, blue, gray, purple, or red.
- Numbness, pain, tingling, or cold feeling of the hands or feet.
- Any sores or wounds on the fingers or toes.
- Muscle pain or weakness, dark urine, or trouble passing urine.
- Erections (hard penis) that happen often or that last a long time.
- Sudden deaths have happened in people with some heart problems or heart defects. Tell the doctor if your child has any heart problem or defect. Tell the doctor if a family member has an abnormal heartbeat or died suddenly. Get medical help right away if your child has any signs of heart problems like chest pain, abnormal heartbeat, or severe dizziness or passing out.
- New or worse behavior and mood changes like change in thinking, anger, and hallucinations have happened with this drug. Tell the doctor if your child or a family member has any mental or mood problems like depression or bipolar illness, or if a family member has committed suicide. Call the doctor right away if your child has hallucinations; change in the way your child acts; or signs of mood changes like depression, thoughts of suicide, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of

interest in life.

- A severe and sometimes deadly problem called serotonin syndrome may happen if your child takes this drug with certain other drugs. Call your child's doctor right away if your child has agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; severe diarrhea, upset stomach, or throwing up; or severe headache.

If your child is or may be sexually active:

- Not able to get or keep an erection.
- Change in sex interest.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Feeling dizzy, tired, or weak.
- Dry mouth.
- Constipation, diarrhea, stomach pain, upset stomach, throwing up, or decreased appetite.
- Trouble sleeping.
- Headache.
- Weight loss.
- Bad taste in your mouth.
- Feeling nervous and excitable.
- Restlessness.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

All products:

- Give early in the day to help prevent sleep problems.
- Give this drug with or without food.

Liquid (suspension):

- Shake well before use.
- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.
- Do not add to food.
- Do not mix with other liquids.
- Put the cap back on after your child is done using a dose.

Oral-disintegrating tablet:

- Do not take this drug out of the blister pack until you are ready to give this drug to your child. Give this drug right away after opening the blister pack. Do not store the removed drug for future use.
- Do not push the tablet out of the foil when opening. Use dry hands to take it from the foil. Place on your child's tongue and let it dissolve. Water is not needed. Do not let your child swallow it whole. Do not let your child chew, break, or crush it.

Extended-release tablets:

- This drug may be chewed or swallowed whole.
- Some strengths of the tablets may have a score line. If needed for your dose, these tablets may be split in half. If splitting the tablets, only split on the score line.

What do I do if my child misses a dose?

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

All products:

- Store at room temperature in a dry place. Do not store in a bathroom.
- Store this drug in a safe place where children cannot see or reach it, and where other people cannot get to it. A locked box or area may help keep this drug safe. Keep all drugs away from pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

Oral-disintegrating tablet:

- Store blister packs in the plastic case that comes with this drug.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's

drug to your child.

- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2025-06-05

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Amphetamine - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering
Cancer Center