

PATIENT & CAREGIVER EDUCATION

Anti-inhibitor Coagulant Complex (Human)

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Feiba

Brand Names: Canada

Feiba NF

Warning

- Blood clots have happened with this drug. Sometimes, blood clots like heart attack and stroke have happened. Tell the doctor if your child has ever had a blood clot. If you have questions, talk with the doctor.
- Call the doctor right away if your child has signs of a blood clot like chest, arm, back, neck, or jaw pain or pressure; coughing up blood; numbness or weakness on 1 side of the body; trouble speaking or thinking; change in balance; change in eyesight; shortness of breath; or swelling, warmth, or pain in the leg or arm.

What is this drug used for?

- It is used to treat or prevent bleeding in people with hemophilia.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has a blood problem called disseminated intravascular coagulation (DIC).

- If your child has had a recent heart attack.
- If your child has a blood clot, has ever had a blood clot, or is at risk of getting a blood clot.
- If your child is taking any of these drugs: Aminocaproic acid or tranexamic acid.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child's blood work and other lab tests checked as you have been told by the doctor.
- Call your child's doctor right away if your child has unexplained bruising or bleeding. This may be a sign that

your child's normal dose is not working as well.

- If your child is also using emicizumab, talk with your child's doctor. Certain blood vessel problems have happened when this drug was used along with emicizumab. Your child will need to be watched closely.
- This drug is made from human plasma (part of the blood) and may have viruses that may cause disease. This drug is screened, tested, and treated to lower the chance that it carries an infection. Talk with the doctor.
- Talk with the doctor before your child travels. Your child will need to bring enough of this drug for use during travel.
- This drug may affect certain lab tests. Be sure the doctor and lab workers know your child uses this drug.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side

effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high or low blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Signs of certain infections (parvovirus B19, hepatitis A) like fever or chills, headache, feeling very sleepy, runny nose, rash, joint pain, tiredness, poor appetite, upset stomach or throwing up, diarrhea, stomach pain, or yellow skin or eyes.
- Feeling very tired or weak.
- Fever or chills.
- Joint pain.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call

your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Diarrhea, upset stomach, or throwing up.
- Feeling dizzy or sleepy.
- Change in taste.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into a vein over a period of time.
- If you will be giving your child the shot, your child's doctor or nurse will teach you how to give the shot.
- Wash your hands before and after use.
- This drug needs to be mixed before use. Follow how to mix as you were told by the doctor.
- When making, do not shake vial.

- Use within 3 hours of making.
- After mixing, do not refrigerate.
- Each vial is for one use only. Throw away any part not used after the dose is given.
- Do not use if the solution is cloudy, leaking, or has particles.
- This drug is colorless to a faint yellow. Do not use if the solution changes color.
- Throw away needles in a needle/sharp disposal box. Do not reuse needles or other items. When the box is full, follow all local rules for getting rid of it. Talk with a doctor or pharmacist if you have any questions.

What do I do if my child misses a dose?

- Call your child's doctor to find out what to do.

How do I store and/or throw out this drug?

- Store at room temperature. Do not freeze.
- Store in the original container to protect from light.
- After mixing, do not refrigerate.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down

a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications,

side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2025-01-22

Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors.
All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Anti-inhibitor Coagulant Complex (Human - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering
Cancer Center