PATIENT & CAREGIVER EDUCATION

Aprepitant

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Aponvie; Cinvanti; Emend; Emend BiPack; Emend TriPack

Brand Names: Canada

Emend; Emend Tri-Pack

What is this drug used for?

- It is used to prevent upset stomach and throwing up.
- It may be given to your child for other reasons. Talk with the doctor.

Injection:

- This drug is not approved for use in children. However, the doctor may decide the benefits of taking this drug outweigh the risks. If your child has been given this drug, ask the doctor for information about the benefits and risks. Talk with the doctor if you have questions about giving this drug to your child.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.

- If your child takes any drugs (prescription or OTC, natural products, vitamins) that must not be taken with this drug, like certain drugs that are used for HIV, infections, seizures, and others. There are many drugs that must not be taken with this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Avoid giving your child grapefruit and grapefruit juice.

If your child is or may be sexually active:

- Birth control pills and other hormone-based birth control may not work as well to prevent pregnancy. Be sure your child uses some other kind of birth control also like a condom when taking this drug and for 1 month after the last dose.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of a very bad skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in the mouth, throat, nose, or eyes.
- Signs of dehydration like dry skin, mouth, or eyes; thirst; fast heartbeat; dizziness; fast breathing; or confusion.
- Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Feeling dizzy, tired, or weak.
- Diarrhea or constipation.
- Decreased appetite.
- Stomach pain or heartburn.
- Hiccups.

- Headache.
- Burping.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

All oral products:

- This drug may be taken with or without food. Be sure to follow the doctor's instructions with regard to food.

Capsules:

- Have your child swallow whole. Do not let your child chew, open, or crush.

Liquid (suspension):

- This drug will be mixed by a doctor or pharmacist.
- When ready to use, place the pre-filled device in the mouth on the inner cheek. Squirt the drug out slowly into the mouth.

What do I do if my child misses a dose?

- Call your child's doctor to find out what to do.

How do I store and/or throw out this drug?

Capsules:

- Store at room temperature in a dry place. Do not store in a bathroom.

Liquid (suspension):

- Store in a refrigerator. Do not freeze.
- Use this drug within 2 days of getting it from the doctor. Throw away any doses left after 2 days.
- When ready to use, this drug may be stored at room temperature for up to 3 hours.

All products:

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2024-03-25

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Aprepitant - Last updated on December 12, 2022
All rights owned and reserved by Memorial Sloan Kettering Cancer Center