

## PATIENT & CAREGIVER EDUCATION

# Asenapine

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

### Brand Names: US

Saphris; Secuado

### Brand Names: Canada

Saphris

### Warning

- There is a higher chance of death in older adults who take this drug for mental problems caused by dementia. Most of the deaths were linked to heart disease or infection. This drug is not approved to treat mental problems caused by dementia.

## **What is this drug used for?**

- It is used to treat bipolar disorder.
- It may be given to your child for other reasons. Talk with the doctor.

## **Skin patch:**

- This drug is not approved for use in children. However, the doctor may decide the benefits of taking this drug outweigh the risks. If your child has been given this drug, ask the doctor for information about the benefits and risks. Talk with the doctor if you have questions about giving this drug to your child.

## **What do I need to tell the doctor BEFORE my child takes this drug?**

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has any of these health problems: Liver problems, low magnesium levels, low potassium levels, or slow heartbeat.
- If your child has ever had a long QT on ECG or other heartbeat that is not normal.
- If your child is taking any drugs that can cause a certain type of heartbeat that is not normal (prolonged QT

interval). There are many drugs that can do this. Ask the doctor or pharmacist if you are not sure.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

### **What are some things I need to know or do while my child takes this drug?**

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child avoid tasks or actions that call for alertness until you see how this drug affects your child. These are things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized vehicles.
- To lower the chance of feeling dizzy or passing out, have your child rise slowly if your child has been sitting or lying

down. Have your child be careful going up and down stairs.

- Dizziness, sleepiness, and feeling less stable may happen with this drug. These may lead to falling, which can cause broken bones or other health problems.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- High blood sugar or diabetes, high cholesterol, and weight gain have happened with drugs like this one. These may raise the chance of heart and brain blood vessel disease.
- Check your child's blood sugar as you have been told by the doctor.
- Alcohol may interact with this drug. Be sure your child does not drink alcohol.
- Talk with your child's doctor before your child uses marijuana, other forms of cannabis, or prescription or OTC drugs that may slow your child's actions.
- Have your child be careful in hot weather or while your child is being active. Have your child drink lots of fluids to stop fluid loss.

## **If your child is pregnant or breast-feeding a baby:**

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.
- Taking this drug in the third trimester of pregnancy may lead to side effects or withdrawal in the newborn.

## **What are some side effects that I need to call my child's doctor about right away?**

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high blood sugar like confusion, feeling sleepy, unusual thirst or hunger, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- Fast, slow, or abnormal heartbeat.

- Shortness of breath.
- Severe dizziness or passing out.
- Trouble controlling body movements, twitching, change in balance, trouble swallowing or speaking.
- Enlarged breasts, nipple discharge, not able to get or keep an erection, or menstrual changes.
- Mouth irritation or mouth sores.
- Seizures.
- Low white blood cell counts have happened with drugs like this one. This may lead to a higher chance of infection. Rarely, infections have been deadly. Tell the doctor if your child has ever had a low white blood cell count. Call the doctor right away if your child has signs of infection like fever, chills, or sore throat.
- A very bad and sometimes deadly health problem called neuroleptic malignant syndrome (NMS) may happen. Call your child's doctor right away if your child has any fever, muscle cramps or stiffness, dizziness, very bad headache, confusion, change in thinking, fast heartbeat, heartbeat that does not feel normal, or is sweating a lot.
- Some people may get a severe muscle problem called tardive dyskinesia. This problem may lessen or go away after stopping this drug, but it may not go away. The risk is greater with diabetes and in older adults, especially

older females. The risk is greater with longer use or higher doses, but it may also occur after short-term use with low doses. Call your child's doctor right away if your child has trouble controlling body movements or problems with the tongue, face, mouth, or jaw like tongue sticking out, puffing cheeks, mouth puckering, or chewing.

### **What are some other side effects of this drug?**

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Feeling dizzy, sleepy, tired, or weak.
- Headache.
- Restlessness.
- Upset stomach or throwing up.
- Stomach pain.
- Numbness or tingling in the mouth.
- Increased appetite.
- Weight gain.
- Trouble sleeping.

- Constipation.
- Change in taste.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

### **How is this drug best given?**

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- Wash and dry your hands before you give this drug to your child. Do not touch the tablet with wet or damp hands.
- Do not take this drug out of the blister pack until you are ready to give this drug to your child. Give this drug right away after opening the blister pack. Do not store the removed drug for future use.
- Place tablet under the tongue and let dissolve.
- Be sure your child does not swallow this drug whole. Do not let your child chew, break, or crush.
- Do not let your child eat or drink for at least 10 minutes



after taking this drug.

- Keep giving this drug to your child as you have been told by your child's doctor or other health care provider, even if your child feels well.

### **What do I do if my child misses a dose?**

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

### **How do I store and/or throw out this drug?**

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

## General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

## Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination

and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

### **Last Reviewed Date**

2025-02-09

### **Copyright**

© 2025 UpToDate, Inc. and its affiliates and/or licensors.

All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

---

Asenapine - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering  
Cancer Center