

## PATIENT & CAREGIVER EDUCATION

# Cannabidiol

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

### Brand Names: US

Epidiolex

### Brand Names: Canada

Epidiolex

### What is this drug used for?

- It is used to help control certain kinds of seizures.

### What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has a feeding tube. This drug is not for use with some types of feeding tubes.

This drug may interact with other drugs or health problems.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

### **What are some things I need to know or do while my child takes this drug?**

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child avoid tasks or actions that call for alertness until you see how this drug affects your child. These are things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized vehicles.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- Do not stop giving this drug to your child all of a sudden without calling the doctor. Your child may have a greater risk of seizures. If your child needs to stop this drug, you will want to slowly stop it as told by the doctor.

- Alcohol may interact with this drug. Be sure your child does not drink alcohol.
- Talk with your child's doctor before your child uses marijuana, other forms of cannabis, or prescription or OTC drugs that may slow your child's actions.
- This drug may affect cannabis (marijuana) lab tests. Tell all of your child's health care providers and lab workers that your child takes this drug.

**If your child is pregnant or breast-feeding a baby:**

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

**What are some side effects that I need to call my child's doctor about right away?**

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever;

wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.

- Signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal.
- Signs of a urinary tract infection (UTI) like blood in the urine, burning or pain when passing urine, feeling the need to pass urine often or right away, fever, lower stomach pain, or pelvic pain.
- Any unexplained bruising or bleeding.
- Trouble walking.
- Like other drugs that may be used for seizures, this drug may rarely raise the risk of suicidal thoughts or actions. The risk may be higher in people who have had suicidal thoughts or actions in the past. Call the doctor right away about any new or worse signs like depression; feeling nervous, restless, or grouchy; panic attacks; or other changes in mood or behavior. Call the doctor right away if any suicidal thoughts or actions occur.

## **What are some other side effects of this drug?**

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Feeling sleepy.
- Decreased appetite.
- Weight loss.
- Diarrhea, upset stomach, or throwing up.
- Feeling tired or weak.
- Trouble sleeping.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

## **How is this drug best given?**

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- Give this drug with or without food but give it the same

way each time. Always give with food or always give on an empty stomach.

- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.
- This drug comes with 2 different sizes of syringe. Be sure you know which size syringe to use to measure the dose. If you are not sure, call the pharmacist.
- Follow how to clean the oral syringes carefully.
- Be sure the oral syringe is dry each time you use it.
- Do not use a household teaspoon or tablespoon to measure this drug. Doing so could lead to the dose being too high.
- If the adapter has not been put in the bottle when you get it, follow how to prepare the bottle as you have been told or read the package insert.
- Keep giving this drug to your child as you have been told by your child's doctor or other health care provider, even if your child feels well.
- Those who have feeding tubes may use this drug. Use as you have been told. Flush the feeding tube after this drug is given.

## What do I do if my child misses a dose?

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

## How do I store and/or throw out this drug?

- Store upright at room temperature. Do not freeze.
- Store in original container.
- Keep lid tightly closed.
- Store in a dry place. Do not store in a bathroom.
- After opening, throw away any part not used after 12 weeks.
- Store this drug in a safe place where children cannot see or reach it, and where other people cannot get to it. A locked box or area may help keep this drug safe. Keep all drugs away from pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

## General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

## Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination



and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

### **Last Reviewed Date**

2025-07-06

### **Copyright**

© 2025 UpToDate, Inc. and its affiliates and/or licensors.

All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

---

Cannabidiol - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering  
Cancer Center