PATIENT & CAREGIVER EDUCATION

Cefotetan

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Cefotan

What is this drug used for?

- It is used to treat or prevent bacterial infections.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If a drug like this one has caused a red blood cell problem called hemolytic anemia.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- If your child has high blood sugar (diabetes) and you test your child's urine glucose, talk with your child's doctor to find out which tests are best to use.
- This drug may affect certain lab tests. Tell all of your child's health care providers and lab workers that your child takes this drug.
- Do not give to your child longer than you have been told. A second infection may happen.
- Alcohol may interact with this drug. Be sure your child does not drink alcohol. Be sure your child does not take

products that have alcohol in them while taking this drug and for at least 72 hours after the last dose. Drinking alcohol or taking products that have alcohol in them, like some cough syrups, may cause flushing, sweating, headaches, and fast heartbeat.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Rarely, some allergic reactions have been deadly.

- Signs of a severe skin reaction (Stevens-Johnson syndrome, toxic epidermal necrolysis, erythema multiforme) like red, swollen, blistered, or peeling skin (with or without fever); skin reaction that looks like rings; red or irritated eyes; or sores in the mouth, throat, nose, or eyes.
- Any unexplained bruising or bleeding.
- Not able to pass urine or change in how much urine is passed.
- Fever, chills, or sore throat.
- Seizures.
- Vaginal itching or discharge.
- Diarrhea is common with antibiotics. Sometimes, diarrhea may be caused by a certain bowel infection (C diff). C diff may happen during treatment or within a few months after the last dose. This infection can be severe or even deadly. Call your child's doctor right away if your child has stomach cramps, fever, long-lasting diarrhea, very loose or watery stools, or bloody stools. Check with your child's doctor before treating diarrhea.
- Rarely, a very bad blood problem called hemolytic anemia has happened with this drug. Sometimes it has been deadly. The chance of getting hemolytic anemia may be higher with this drug than with other drugs like

this one. Call your doctor right away if your child feels very tired or weak, or develops dark urine or yellow skin or eyes while using this drug or within 3 weeks after stopping it.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Diarrhea.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into a muscle or as an infusion into a vein over a period of time.

What do I do if my child misses a dose?

- Call your child's doctor to find out what to do.

How do I store and/or throw out this drug?

- If you need to store this drug at home, talk with your child's doctor, nurse, or pharmacist about how to store it.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.

- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the

Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2024-05-08

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Cefotetan - Last updated on December 12, 2022
All rights owned and reserved by Memorial Sloan Kettering Cancer Center

Memorial Sloan Kettering Cancer Center