

Chloramphenicol (Systemic)

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: Canada

Chloromycetin Succinate

Warning

- Very bad and sometimes deadly blood problems have happened with this drug. This may cause more chance of getting an infection, bleeding problems, or slow healing. Aplastic anemia that happened with this drug has led to a certain kind of cancer (leukemia). Blood problems have happened after both short-term use and long-term use. Have your child's blood work checked while taking this drug. Talk with the doctor.

What is this drug used for?

- It is used to treat bacterial infections.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.

- If your child has any of these health problems: Bone marrow disease or low blood cell counts.
- If your child is taking any drugs that can raise the chance of blood problems. There are many drugs that can do this. Ask your doctor or pharmacist if you are not sure.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Do not give to your child longer than you have been told. A second infection may happen.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered,

or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.

- Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal.
- Signs of bleeding like throwing up or coughing up blood; vomit that looks like coffee grounds; blood in the urine; black, red, or tarry stools; bleeding from the gums; abnormal vaginal bleeding; bruises without a cause or that get bigger; or bleeding you cannot stop.
- Feeling very tired or weak.
- Feeling confused.
- Depression or other mood changes.
- Change in eyesight.
- Diarrhea is common with antibiotics. Rarely, a severe form called C diff–associated diarrhea (CDAD) may happen. Sometimes, this has led to a deadly bowel problem. CDAD may happen during or a few months after taking antibiotics. Call your child’s doctor right away if your child has stomach pain, cramps, or very loose, watery, or bloody stools. Check with your child’s doctor before treating diarrhea.
- Very bad and sometimes deadly reactions have happened in premature and newborn babies. This has happened in a baby born to a mother who got this drug during labor. It has also happened in a 3-month old. Most of the time, this drug was used within the first 48 hours of life. Signs first show up after 3 to 4 days of getting this drug. Call your child’s doctor right away if your child has swelling of the stomach with or without throwing up, blue or gray skin color, or trouble breathing.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Headache.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- It is given into a vein for a period of time.

What do I do if my child misses a dose?

- Call your child's doctor to find out what to do.

How do I store and/or throw out this drug?

- If you need to store this drug at home, talk with your child's doctor, nurse, or pharmacist about how to store it.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and

pets.

- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2023-07-12

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Chloramphenicol (Systemic - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering
Cancer Center