PATIENT & CAREGIVER EDUCATION

Cromolyn (Systemic

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Gastrocrom

Brand Names: Canada

Nalcrom

What is this drug used for?

- It is used to treat mastocytosis.
- It is used to treat food allergies.
- It may be given to your child for other reasons. Talk with the doctor.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor

about the allergy and what signs your child had.

This drug may interact with other drugs or health problems.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

All products:

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

Capsules:

- If your child has had a life-threatening reaction to foods before, talk with the doctor. You may need to give your child other drugs with this drug.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

All products:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.

Capsules:

- Coughing.
- Muscle pain or weakness.
- Shortness of breath, a big weight gain, or swelling in the arms or legs.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

Capsules:

- Diarrhea, upset stomach, or throwing up.
- Stomach pain.
- Headache.
- Trouble sleeping.
- Sneezing.
- Bad taste in your mouth.
- Joint pain.

Liquid (concentrate):

- Diarrhea.
- Headache.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

Capsules:

- Give 15 to 20 minutes before meals.
- Your child may swallow whole or you may mix the contents of the capsule with water. It is best to mix this drug with water before giving it to your child. If mixing it with water, mix as you have been told by the doctor. Have your child take the mixture right away. Do not store it for later use.
- Keep giving this drug to your child as you have been told by your child's doctor or other health care provider, even if your child feels well.
- Do not stop giving this drug to your child all of a sudden without calling the doctor. Your child may have a greater risk of side effects. If your child needs to stop this drug, you will want to slowly stop it as told by the doctor.

Liquid (concentrate):

- Give by mouth only. This product is not to be used for breathing in with a nebulizer.
- Do not use if the solution is cloudy, leaking, or has particles.
- Do not use if solution changes color.
- Give 30 minutes before meals and at bedtime unless your child's doctor has told you otherwise.
- Do not let your child swallow the ampul. Open it and empty the liquid into a glass of water. Mix and have your child drink.
- Keep giving this drug to your child as you have been told by your child's doctor or other health care provider, even if your child feels well.

What do I do if my child misses a dose?

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

Capsules:

- Store at room temperature in a dry place. Do not store in a bathroom.

Liquid (concentrate):

- Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom.
- Store unused containers in foil pouch until use.

All products:

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information

leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.

- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a

specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date
2025-09-16

Copyright
© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Cromolyn (Systemic - Last updated on December 12, 2022
All rights owned and reserved by Memorial Sloan Kettering Cancer Center