PATIENT & CAREGIVER EDUCATION

Cyclopentolate and Phenylephrine

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Cyclomydril

What is this drug used for?

- It makes the eye pupils larger.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has narrow-angle glaucoma.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's

drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

For all patients taking this drug:

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child avoid tasks or actions that call for clear eyesight while your child's pupils are dilated.
- Bright lights may bother your child. Have your child wear sunglasses.
- Severe heart problems like abnormal heartbeats and heart attack have happened with the 10% strength of phenylephrine. Most of the time, these heart problems happened in people who had heart disease. Sometimes, they were deadly. If you have questions, talk with the doctor.
- If the patient is a child, use this drug with care. The risk of some side effects may be higher in children.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

Infants:

- Watch infants closely for at least 30 minutes after giving this drug.
- If this drug is given to an infant, do not feed the infant after this drug is given. Talk with the doctor to find out when the infant can be fed.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

For all patients taking this drug:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever;

wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.

- Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Chest pain or pressure.
- Fast, slow, or abnormal heartbeat.
- Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight.
- Change in eyesight, eye pain, or severe eye irritation.
- Flushing.
- Trouble passing urine.
- Feeling confused.
- Behavior problems.
- Hallucinations (seeing or hearing things that are not there).
- Seizures.
- Change in balance.
- Trouble speaking.
- Restlessness.
- Feeling sleepy.

- Dry mouth, nose, or throat.

Newborns:

- Constipation; diarrhea; dark or bloody stools; swollen or tender belly; red or gray belly; trouble feeding; green vomit; trouble breathing; sluggishness; slow heart rate; or temperature changes.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Eye irritation.
- Burning.
- Blurred eyesight.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- For the eye only.

What do I do if my child misses a dose?

- Call your child's doctor to find out what to do.

How do I store and/or throw out this drug?

- If you need to store this drug at home, talk with your child's doctor, nurse, or pharmacist about how to store it.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or

medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2025-05-30

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Cyclopentolate and Phenylephrine - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center

Memorial Sloan Kettering Cancer Center