

Doravirine

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Pifeltro

Brand Names: Canada

Pifeltro

What is this drug used for?

- It is used to treat HIV infection.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child weighs less than 77 pounds (35 kilograms).
- If your child is taking any of these drugs or has taken any of these drugs in the last 4 weeks: Carbamazepine, enzalutamide, mitotane, oxcarbazepine, phenobarbital, phenytoin, rifampin, rifapentine, or St. John's wort.
- If your child is taking any of these drugs: Efavirenz, etravirine, or nevirapine.

If your child is breast-feeding a baby:

- Be sure your child does not breast-feed a baby while taking this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Do not run out of this drug.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- This drug is not a cure for HIV. Keep giving this drug to your child as you have been told by the doctor or other health care provider, even if your child feels well. Be sure your child stays under the care of the doctor and has regular appointments.
- This drug does not stop the spread of diseases like HIV or hepatitis that are passed through blood. Be sure needles and other things like toothbrushes or razors are not shared.

If your child is or may be sexually active:

- This drug does not stop the spread of diseases like HIV or hepatitis that are passed through having sex. Be sure your child does not have any kind of sex without using a latex or polyurethane condom.

If your child is pregnant:

- Tell the doctor if your child is pregnant or becomes pregnant. You will need to talk about the benefits and risks of your child using this drug while pregnant.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Changes in your child's immune system can happen when your child starts taking drugs to treat HIV. If your child has an infection that you did not know was there, it may show up when your child takes this drug. Tell your child's doctor right away if your child has any new signs after starting this drug, even after taking it for several months. This includes signs of infection like fever, sore throat, weakness, cough, or shortness of breath.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Upset stomach.
- Feeling dizzy, tired, or weak.
- Strange or odd dreams.

- Stomach pain or diarrhea.
- Headache.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- Give this drug with or without food.
- Give this drug at the same time of day.
- Keep giving this drug to your child as you have been told by your child's doctor or other health care provider, even if your child feels well.
- It is important that your child does not miss or skip a dose of this drug during treatment.

What do I do if my child misses a dose?

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.
- If you are not sure what to do if your child misses a dose, call the doctor.

How do I store and/or throw out this drug?

- Store tablets in the original container at room temperature. Keep the cap tightly closed.
- Store in a dry place. Do not store in a bathroom.

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient.

UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2023-12-22

Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your care team directly. If you're a patient at MSK and you need to reach a provider after 5 p.m., during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Doravirine - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering
Cancer Center