

## PATIENT & CAREGIVER EDUCATION

# Fluoride

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

### Brand Names: US

Clinpro 5000; Denta 5000 Plus; Denta 5000 Plus Sensitive; DentaGel; Easygel; Easygel [OTC]; Fluoridex; Fluoridex Daily Renewal; Fluoridex Enhanced Whitening; Fluoridex Sensitivity Relief; FluoriMax 5000; FluoriMax 5000 Sensitive; Fluoritab [DSC]; Fraiche 5000 Dental; Fraiche Rinse [OTC] [DSC]; Gel-Kam [OTC]; Just for Kids [OTC]; Just Right 5000; NaFrinse Daily Acidulated [DSC]; NaFrinse Daily/Neutral [DSC]; NaFrinse Drops [DSC]; NaFrinse Weekly [DSC]; NaFrinse [DSC]; Omni Gel [OTC]; Parodontax [OTC]; PerioMed [OTC]; PreviDent; PreviDent 5000 Booster Plus; PreviDent 5000 Dry Mouth; PreviDent 5000 Kids; PreviDent 5000 Ortho Defense; PreviDent 5000 Plus; Sensodyne Complete Protection [OTC]; Sensodyne Pronamel Kids [OTC]; Sensodyne Rapid Relief [OTC]; Sensodyne Repair & Protect [OTC]; Sensodyne

**Sensitivity & Gum [OTC]; Sensodyne Sensitivity Gum-Enam [OTC]; SF; SF 5000 Plus; Sodium Fluoride 5000 Enamel; Sodium Fluoride 5000 Plus; Sodium Fluoride 5000 PPM; Sodium Fluoride 5000 Sensitive**

**Brand Names: Canada**  
**Butler Prophy Paste**

**What is this drug used for?**

**All products:**

- This supplement is used to prevent cavities.

**Sodium fluoride and potassium nitrate:**

- It is used to help sensitive teeth.

**What do I need to tell the doctor BEFORE my child takes this drug?**

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has any of these health problems: Bone problems, joint problems, kidney problems, or ulcers.
- If the fluoride in your drinking water is greater than or equal to 0.6 parts per million.

**This is not a list of all drugs or health problems that interact with this drug.**

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

## **What are some things I need to know or do while my child takes this drug?**

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Tell the dentist if your child's teeth become stained or spotted.
- Take care of your child's teeth. See a dentist often.
- This drug may cause harm if a large amount is swallowed. If a large amount of this drug is swallowed, call a doctor or poison control center right away.
- Different brands of this drug may be for use in different ages of children. Talk with the doctor before giving this drug to a child.

## If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

## What are some side effects that I need to call my child's doctor about right away?

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Severe upset stomach or throwing up.

## What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if your child has any side effects that bother your child or do not go away.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

### **How is this drug best given?**

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

#### **Chewable tablets, liquid (drops), or tablets:**

- Give this drug with or without food. Give with food if it causes an upset stomach.
- Give this drug before bedtime, after you brush your child's teeth, unless the doctor tells you otherwise.
- Some products are used every day while others are used 1 time a week. Follow what you have been told by your doctor.
- Do not give antacids that have magnesium or aluminum at the same time as this drug. Talk with the doctor or pharmacist.
- Do not give products that have calcium like dairy products within 2 hours of this drug.

## **Liquid (drops):**

- Measure liquid doses carefully. Use the measuring device that comes with this drug.
- Some brands of the drops need to be mixed with water or fruit juice. Check to see how your brand is to be taken.

## **Chewable tablets:**

- Have your child chew well before swallowing.
- Some brands of this drug may also be dissolved in your child's mouth. Check with the doctor or pharmacist.

## **Oral cream and paste:**

- Use in place of your child's normal toothpaste, unless the doctor says otherwise.
- Have your child brush with this drug for 2 minutes and then spit out.
- Do not let your child swallow this drug.
- Make sure your child does not eat, drink, or rinse for at least 30 minutes after using this drug.

## **Sodium fluoride and potassium nitrate:**

- Have your child rinse well and spit out rinse water.

## **Gel:**

- Use in place of your child's normal toothpaste, unless the doctor says otherwise.
- Apply a pea-sized amount and have your child brush well. Let the drug stay on your child's teeth for 1 minute before spitting out.
- Do not let your child swallow this drug.
- Make sure your child does not eat, drink, or rinse for at least 30 minutes after using this drug.

## **Oral rinse:**

- Have your child swish it in the mouth as long as you were told and then spit out.
- Do not let your child swallow this drug.
- Make sure your child does not eat, drink, or rinse for at least 30 minutes after using this drug.
- Some products need to be mixed before using. If the product needs to be mixed, do not mix until you are ready to use it. Use right away after mixing.

## **What do I do if my child misses a dose?**

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip

the missed dose and go back to your child's normal time.

- Do not give 2 doses at the same time or extra doses.

## How do I store and/or throw out this drug?

### All products:

- Store at room temperature.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

### Liquid (drops):

- Protect from light.

### General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other

health care provider.

- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

## Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the

use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

## Last Reviewed Date

2024-07-12

## Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

Fluoride - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering  
Cancer Center

