

PATIENT & CAREGIVER EDUCATION

Isavuconazonium Sulfate

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Cresemba

Brand Names: Canada

Cresemba

What is this drug used for?

- It is used to treat fungal infections.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has Familial Short QT syndrome.
- If your child takes any drugs (prescription or OTC, natural products, vitamins) that must not be taken with this drug. This includes rifampin, St. John's wort, or certain drugs used for seizures. There are many drugs that must not be taken with this drug.

If your child is breast-feeding a baby:

- Be sure your child does not breast-feed a baby while taking this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child's blood work and other lab tests checked as you have been told by the doctor.

If your child is or may be sexually active:

- This drug may cause harm to the unborn baby if your child takes it during pregnancy.
- If your child may become pregnant, have your child use birth control while taking this drug. Your child will also need to use birth control for 28 days after the last dose unless you have been told otherwise. Be sure you know how long your child needs to use birth control. If your child gets pregnant, call the doctor right away.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs

or symptoms that may be related to a very bad side effect:

All products:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat. Rarely, some allergic reactions have been deadly.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Signs of electrolyte problems like mood changes; confusion; muscle pain, cramps, or spasms; weakness; shakiness; change in balance; an abnormal heartbeat; seizures; loss of appetite; or severe upset stomach or throwing up.
- Chest pain or pressure, a fast heartbeat, or an abnormal heartbeat.
- Feeling confused.
- Shortness of breath.
- Swelling in the arms or legs.
- Dizziness or passing out.
- Flushing.
- Sweating a lot.
- Liver problems have happened with this drug. Sometimes, liver problems have been deadly in people with severe health problems. Call the doctor right away if your child has signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- A severe skin reaction (Stevens-Johnson syndrome) may happen. It can cause severe health problems that may not go away, and sometimes death. Get medical help right away if your child has signs like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in the mouth, throat, nose, or eyes.

Injection:

- Some patients have had very bad side effects during the infusion. Tell the doctor if your child has a burning, numbness, or tingling feeling that is not normal; chills; dizziness or passing out; shortness of breath; or any other bad effects during the infusion.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

All products:

- Back pain.
- Constipation, diarrhea, stomach pain, upset stomach, throwing up, or decreased appetite.
- Cough.
- Feeling tired or weak.
- Headache.
- Trouble sleeping.
- Anxiety.

Injection:

- Irritation where this drug is given.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

Capsules:

- Give this drug with or without food.
- Have your child swallow whole. Do not let your child chew, crush, open, or dissolve.
- Do not take this drug out of the blister pack until you are ready to give this drug to your child. Give this drug right away after opening the blister pack. Do not store the removed drug for future use.
- Be sure to only open the capsule pocket. Do not open the pocket that contains the desiccant (protects the capsule from moisture). Do not swallow or use the desiccant.
- Use as you have been told, even if your child's signs get better.
- Do not put this form of this drug down a feeding tube.

Injection:

- It is given as an infusion into a vein over a period of time.
- If needed, this form of this drug may be given through a feeding tube instead of as an injection. This drug must be used within 1 hour if mixed to be given through a feeding tube.

What do I do if my child misses a dose?

Capsules:

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

Injection:

- Call your child's doctor to find out what to do.

How do I store and/or throw out this drug?

Capsules:

- Store in the original container at room temperature.
- Store in a dry place. Do not store in a bathroom.

Injection:

- If you need to store this drug at home, talk with your child's doctor, nurse, or pharmacist about how to store it.

All products:

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how

much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2025-05-16

Copyright

© 2025 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Isavuconazonium Sulfate - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering
Cancer Center