PATIENT & CAREGIVER EDUCATION

Lisdexamfetamine

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Arynta; Vyvanse

Brand Names: Canada

AG-Lisdexamfetamine; APO-Lisdexamfetamine; JAMP-Lisdexamfetamine; PRO-Lisdexamfetamine; SANDOZ Lisdexamfetamine; TARO-Lisdexamfetamine; TEVA Lisdexamfetamine; Vyvanse

Warning

- This drug has a high risk of misuse. This can lead to alcohol or drug use disorder. Misuse or abuse of this drug can lead to overdose or death. The risk is higher with higher doses or when used in ways that are not approved like snorting or injecting it. Do not give this drug to anyone else. This drug may also be habit-forming if taken for a long time. Do not give for longer than you have been told by the doctor. Give only as you were told. Tell the doctor if your child has ever had alcohol or drug use disorder. Your child will be watched closely while taking this drug. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs as you have been told.

What is this drug used for?

- It is used to treat attention deficit problems with hyperactivity.
- It is used to treat binge eating disorder.
- It may be given to your child for other reasons. Talk with the doctor.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child or a family member has any of these health problems: Blood vessel disease, high blood pressure, heart structure problems or other heart problems, or Tourette's syndrome or tics.
- If your child has any of these health problems: Glaucoma, agitation, anxiety, or overactive thyroid.
- If your child has ever had a stroke.
- If your child is taking any of these drugs: Acetazolamide or sodium bicarbonate.
- If your child has taken isocarboxazid, linezolid, methylene blue, phenelzine, tranylcypromine, selegiline, or rasagiline in the last 14 days. Severe high blood pressure may happen.

If your child is breast-feeding a baby:

- Be sure your child does not breast-feed a baby while taking this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes

without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- This drug is not approved for weight loss. If you have questions, talk with the doctor.
- Have your child avoid tasks or actions that call for alertness or clear eyesight until you see how this drug affects your child. These are things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized vehicles.
- This drug may cause high blood pressure.
- Have your child's blood pressure and heart rate checked often.
- Your child may need to have some heart tests before starting this drug. If you have questions, talk with your child's doctor.
- This drug may affect radioactive imaging tests and certain other lab tests. Tell all of your child's health care providers and lab workers that your child takes this drug.
- Talk with the health care provider before giving OTC products that may increase blood pressure. These include cough or cold remedies, diet pills, stimulants, ibuprofen or like products, and certain natural products or supplements.
- If your child has been taking this drug for a long time or at high doses, it may not work as well and your child may need higher doses to get the same effect. This is known as tolerance. Call the doctor if this drug stops working well. Do not give more than ordered.
- Long-term or regular use of this drug may lead to dependence. Stopping this drug all of a sudden may lead to signs of withdrawal. Talk to your child's doctor before you lower the dose or stop giving this drug. You will need to follow the doctor's instructions. Tell your child's doctor if your child has any

bad effects.

- This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor.
- The risk of weight loss and other side effects is higher in children younger than 6 years of age. This drug is not for use in children younger than 6 years of age.

If your child is pregnant:

- Tell the doctor if your child is pregnant or becomes pregnant. You will need to talk about the benefits and risks of your child using this drug while pregnant.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Fast or abnormal heartbeat.
- Trouble controlling body movements.
- Trouble controlling sounds, like humming, throat clearing, yelling, or making loud noises.
- Change in color of hands, feet, or other areas. Skin may turn pale, blue, gray, purple, or red.
- Numbness, pain, tingling, or cold feeling of the hands or feet.

- Any sores or wounds on the fingers or toes.
- Muscle pain or weakness, dark urine, or trouble passing urine.
- Sudden deaths have happened in people with some heart problems or heart defects. Tell the doctor if your child has any heart problem or defect. Tell the doctor if a family member has an abnormal heartbeat or died suddenly. Get medical help right away if your child has any signs of heart problems like chest pain, abnormal heartbeat, or severe dizziness or passing out.
- New or worse behavior and mood changes like change in thinking, anger, and hallucinations have happened with this drug. Tell the doctor if your child or a family member has any mental or mood problems like depression or bipolar illness, or if a family member has committed suicide. Call the doctor right away if your child has hallucinations; change in the way your child acts; or signs of mood changes like depression, thoughts of suicide, nervousness, emotional ups and downs, thinking that is not normal, anxiety, or lack of interest in life.
- A severe skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) may happen. It can cause severe health problems that may not go away, and sometimes death. Get medical help right away if your child has signs like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in the mouth, throat, nose, or eyes.
- A severe and sometimes deadly problem called serotonin syndrome may happen if your child takes this drug with certain other drugs. Call your child's doctor right away if your child has agitation; change in balance; confusion; hallucinations; fever; fast or abnormal heartbeat; flushing; muscle twitching or stiffness; seizures; shivering or shaking; sweating a lot; severe diarrhea, upset stomach, or throwing up; or severe headache.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Constipation, diarrhea, stomach pain, upset stomach, throwing up, or decreased appetite.
- Weight loss.
- Trouble sleeping.
- Anxiety.
- Dry mouth.
- Dizziness.
- Feeling jittery.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

All products:

- Give this drug at the same time of day.
- Give this drug with or without food.
- Give in the morning. Giving this drug in the afternoon or later may cause trouble sleeping.

Capsules:

- Have your child swallow whole. Do not let your child chew, break, or crush.
- You may open the capsule and mix the powder with water, orange juice, or yogurt. If needed, a spoon may be used to break apart powder that is stuck together. Mix well.
- If mixed, have your child swallow the mixed drug right away. Do not store for

use at a later time.

Chewable tablets:

- Have your child chew well before swallowing.

Oral solution:

- If the adapter has not been put in the bottle when you get it, prepare the bottle as you have been told or read the package insert. Use a new adapter with each new bottle. Do not take the adapter out of the bottle after it has been put in.
- Measure liquid doses carefully. Use the measuring device that comes with this drug.
- Have your child sit upright to take a dose. Place the tip of the syringe into the mouth on the inside of the cheek. Gently push the plunger until all the drug is gone. Do not squirt quickly. This could cause choking.

What do I do if my child misses a dose?

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

All products:

- Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom.
- Store this drug in a safe place where children cannot see or reach it, and where other people cannot get to it. A locked box or area may help keep this drug safe. Keep all drugs away from pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you

have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

Oral solution:

- Throw away any drug that is left 30 days after opening.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of

this information is governed by the Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2026-05-21

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Lisdexamfetamine - Last updated on December 12, 2022
All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering Cancer Center