

Mecasermin

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Increlex

Brand Names: Canada

Increlex

What is this drug used for?

- It is used to help growth in certain children.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has or has ever had cancer, has a health problem that raises the risk of cancer, or has a family member who has had cancer.
- If your child's bones are no longer growing (closed epiphyses).
- If the patient is a newborn or infant. This drug has benzyl alcohol in it. If possible, avoid products with benzyl alcohol in newborns or infants. Serious side effects can happen in these children with some doses of benzyl alcohol, including if given with other drugs that have benzyl alcohol.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child avoid tasks or actions that call for alertness for 2 to 3 hours after getting this drug and until you see how this drug affects your child. This includes things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized vehicles.
- If your child has high blood sugar (diabetes), talk with the doctor. This drug may lower blood sugar. High blood sugar drugs may need to be changed.
- Check your child's blood sugar as you have been told by the doctor.
- Have your child's blood work and other lab tests checked as you have been told by the doctor.
- Get your child an eye exam as you have been told by the doctor.
- Monitor growth carefully.
- If your child has an abnormal curve in the spine (scoliosis), talk with your doctor. This drug could make it worse in children who are still growing.
- This drug may cause high cholesterol and triglyceride levels. Talk with the doctor.
- Cases of cancer have happened in children taking this drug. It is not known if this drug caused this effect. If you have questions, talk with your child's doctor.
- Allergic reactions have happened with this drug. Sometimes, these have

been very bad and people have had to go to the hospital. Talk with the doctor.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Dizziness or passing out.
- Pale skin.
- Clammy skin.
- Change in eyesight.
- Snoring.
- Breathing problems during sleep (sleep apnea).
- Larger tonsils.
- Ear pain.
- Change in hearing.
- Seizures.
- Thick skin, pits, or lumps where the injection was given.

- A heartbeat that does not feel normal.
- Low blood sugar may happen. Signs may be dizziness or passing out, blurred eyesight, mood changes, slurred speech, headache, feeling sleepy or weak, shaking, fast heartbeat, confusion, hunger, sweating, or seizures. Call the doctor right away if any of these signs happen. Follow what you have been told to do if low blood sugar happens. This may include taking glucose tablets, liquid glucose, or some fruit juices.
- Raised pressure in the brain has happened with this drug. Call your child's doctor right away if your child has a headache; dizziness; upset stomach; ringing in the ears; eye pain; or eyesight problems like blurred eyesight, seeing double, or loss of eyesight.
- Children who use this drug may rarely have a bone problem in the hip (slipped growth plate). This can also lead to further bone injury. Call the doctor right away if your child has hip or knee pain or a limp.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Joint pain.
- Bruising where the shot is given.
- Irritation where the shot is given.
- Pain in arms or legs.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into the fatty part of the skin in the upper arm, thigh, buttocks, or stomach area.
- If you will be giving your child the shot, your child's doctor or nurse will teach you how to give the shot.
- This drug must be given within 20 minutes before or after your child's meal or snack.
- If your child is not able to eat, do not give a dose and do not make up for the missed dose.
- Wash your hands before and after use.
- Move the site where you give the shot with each shot.
- Do not use if the solution is cloudy, leaking, or has particles.
- This drug is clear and colorless. Do not use if the solution changes color.
- Throw away needles in a needle/sharp disposal box. Do not reuse needles or other items. When the box is full, follow all local rules for getting rid of it. Talk with a doctor or pharmacist if you have any questions.

What do I do if my child misses a dose?

- Skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store in a refrigerator. Do not freeze.
- Protect from light.
- Do not use if it has been frozen.
- After opening, throw away any part not used after 30 days.

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any

warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2025-07-30

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Mecasermin - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center

Last Updated

December 12, 2022

Learn about our [Health Information Policy](#).



Memorial Sloan Kettering
Cancer Center