

PATIENT & CAREGIVER EDUCATION

Mometasone (Oral Inhalation)

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Asmanex (120 Metered Doses); Asmanex (14 Metered Doses); Asmanex (30 Metered Doses); Asmanex (60 Metered Doses); Asmanex (7 Metered Doses) [DSC]; Asmanex HFA

Brand Names: Canada

Asmanex Twisthaler

What is this drug used for?

- It is used to treat asthma.
- Do not give this drug to treat an asthma attack. Use a rescue inhaler. Talk with the doctor.

What do I need to tell the doctor BEFORE my child takes this drug?

All products:

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.

Inhaler (powder):

- If your child is allergic to milk, talk with the doctor.

This drug may interact with other drugs or health problems.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's

doctors, nurses, pharmacists, and dentists.

- Call the doctor right away if your child has breathing problems that get worse, if the rescue inhaler does not work as well, or if your child needs to use the rescue inhaler more often.
- It may take a few weeks to see the full effect.
- If your child has been taking this drug for many weeks, talk with your child's doctor before stopping. You may want to slowly stop this drug.
- Tell the doctor if your child has missed a dose or recently stopped this drug and feels very tired, weak, or shaky, or has a fast heartbeat, confusion, sweating, or dizziness.
- Your child may have more chance of getting infections. Have your child wash hands often. Avoid crowds and people with infections, colds, or flu.
- Chickenpox and measles can be very bad or even deadly in some people taking steroid drugs like this drug. Avoid having your child near anyone with chickenpox or measles if your child has not had these health problems before. If your child has been exposed to chickenpox or measles, talk with the doctor.
- When changing from an oral steroid to another form

of steroid, there may be severe and sometimes deadly side effects. Signs like weakness, feeling tired, dizziness, upset stomach, throwing up, confusion, or low blood sugar may happen. Call the doctor right away if your child has any of these signs. If your child has a bad injury, has surgery, or any type of infection, they may need extra doses of oral steroids. These extra steroids will help your child's body deal with these stresses. Have your child wear medical alert ID (identification).

- Long-term use may raise the chance of cataracts, glaucoma, or weak bones (osteoporosis). Talk with your child's doctor.
- Your child may need to have a bone density test. Talk with the doctor.
- Have your child's eye pressure checked if your child is on this drug for a long time. Talk with the doctor.
- If the patient is a child, use this drug with care. The risk of some side effects may be higher in children.
- This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

All products:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal.

- Signs of a weak adrenal gland like a severe upset stomach or throwing up, severe dizziness or passing out, muscle weakness, feeling very tired, mood changes, decreased appetite, or weight loss.
- Redness or white patches in mouth or throat.
- Bone or joint pain.
- Change in eyesight.
- Flushing.
- This drug can cause very bad breathing problems right after your child takes a dose. Sometimes, this may be life-threatening. If your child has trouble breathing, breathing that is worse, wheezing, or coughing after using this drug, have your child use a rescue inhaler and get medical help right away.

Inhaler (aerosol):

- Flu-like signs.

Inhaler (powder):

If your child has menstrual periods:

- Period (menstrual) pain.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call

your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

All products:

- Headache.
- Nose or throat irritation.

Inhaler (powder):

- Stomach pain.
- Upset stomach.
- Muscle pain.
- Back pain.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

All products:

- For breathing in only.
- Keep giving this drug to your child as you have been told by your child's doctor or other health care provider, even if your child feels well.
- Have your child rinse out the mouth after each use. Do not let your child swallow the rinse water. Have your child spit it out.
- Some inhalers have a dose counter to keep track of how many doses are left. If your inhaler has a dose counter, throw the inhaler away when the dose counter reaches "0."

Inhaler (aerosol):

- Prime before first use or if not used for more than 5 days. To prime the inhaler, spray 4 times into the air away from the face and eyes. Shake well before each spray.
- Shake well before use.
- Put the cap back on after your child is done using a dose.
- Do not use near an open flame or while smoking. It may burst.
- Only use the device that comes with this drug. Do not

use any other devices.

- Do not take the device apart or wash it in water. Clean the mouthpiece at least every 7 days of use by using a dry wipe.
- Keep the spray away from the eyes. If this drug gets in the eyes, rinse right away with water. If eye irritation or redness lasts, call the doctor.

Inhaler (powder):

- Clean mouthpiece by wiping with a dry tissue or cloth. Do not wash or put in water.
- Do not let your child breathe out into the inhaler. Put the cap back on after giving a dose.

What do I do if my child misses a dose?

- Skip the missed dose and go back to your child's normal time unless your child's doctor tells you to do something else.
- Do not give 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

All products:

- Store at room temperature in a dry place. Do not store in a bathroom.

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

Inhaler (aerosol):

- Protect from heat or open flame. Do not puncture or burn even if it seems empty.

Inhaler (powder):

- Throw away inhaler after 45 days of opening the pouch, when it says 0, or locks you from using it.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse,

pharmacist, or other health care provider.

- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability

relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2022-06-20

Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Mometasone (Oral Inhalation - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center

