

PATIENT & CAREGIVER EDUCATION

Nelarabine

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Arranon

Brand Names: Canada

Atriance

Warning

- This drug may cause nervous system problems. Sometimes, nervous system problems have not gone away even after treatment with this drug was stopped. Some nervous system problems have been severe, life-threatening, or even deadly. Call your child's doctor right away if your child feels very sleepy, very dizzy, confused, or is not able to move. Call your child's doctor right away if your child has a burning, numbness, or tingling feeling that is not normal; a change in balance; problems with daily living like buttoning clothes; seizures; tripping when walking; or weakness.

What is this drug used for?

- It is used to treat types of leukemia and lymphoma.

What do I need to tell the doctor **BEFORE** my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child is taking pentostatin.

If your child is breast-feeding a baby:

- Be sure your child does not breast-feed a baby while taking this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Your child may feel sleepy during treatment and for several days after treatment. Have your child avoid tasks or actions that call for alertness until your child does not feel sleepy anymore. These are things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized

vehicles.

- This drug may lower the ability of the bone marrow to make blood cells that the body needs. If blood cell counts get very low, this can lead to bleeding problems, infections, or anemia. If you have questions, talk with the doctor.
- Your child may bleed more easily. Make sure your child is careful and avoids injury. Be sure your child has a soft toothbrush.
- Your child may have more chance of getting infections. Have your child wash hands often. Avoid crowds and people with infections, colds, or flu.
- If your child has high blood sugar (diabetes), you will need to watch your child's blood sugar closely.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- Talk with the doctor before your child gets any vaccines. Use of some vaccines with this drug may either raise the chance of very bad infection or make the vaccine not work as well.
- If the patient is a child, use this drug with care. The risk of some side effects may be higher in children.
- If your child is of childbearing age, a pregnancy test will need to be done before starting this drug to make sure

your child is not pregnant.

If your child is or may be sexually active:

- This drug may cause harm to an unborn baby. Your child must use birth control while taking this drug. If your child gets pregnant, call your child's doctor right away.
- If your child's sex partner may become pregnant, have them use a condom while taking this drug and for 3 months after the last dose. If your child's partner gets pregnant, call the doctor right away.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color

of sputum, pain with passing urine, mouth sores, or wound that will not heal.

- Signs of bleeding like throwing up or coughing up blood; vomit that looks like coffee grounds; blood in the urine; black, red, or tarry stools; bleeding from the gums; abnormal vaginal bleeding; bruises without a cause or that get bigger; or bleeding you cannot stop.
- Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, fast or abnormal heartbeat, severe dizziness or passing out, increased thirst, seizures, feeling very tired or weak, decreased appetite, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or severe upset stomach or throwing up.
- Signs of high blood sugar like confusion, feeling sleepy, unusual thirst or hunger, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- Shortness of breath.
- Pale skin.
- Chest pain or pressure, a fast heartbeat, or passing out.
- Swelling.
- Depression.
- Patients with cancer who take this drug may be at

greater risk of getting a severe health problem called tumor lysis syndrome (TLS). This may lead to death. Call your child's doctor right away if your child has a fast or abnormal heartbeat; any passing out; trouble passing urine; muscle weakness or cramps; upset stomach, throwing up, diarrhea, or not able to eat; or feels sluggish.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Feeling dizzy, sleepy, tired, or weak.
- Headache.
- Back, muscle, or joint pain.
- Mouth irritation or mouth sores.
- Swelling of belly.
- Trouble sleeping.
- Pain in arms or legs.
- Shakiness.
- Nosebleed.

- Constipation.
- Stomach pain.
- Diarrhea, throwing up, upset stomach, and decreased appetite are common with this drug. If these happen, talk with your child's doctor about ways to lower these side effects. Call your child's doctor right away if any of these effects bother your child, do not get better, or are severe.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- It is given as an infusion into a vein over a period of time.
- Other drugs will be given with this drug to help avoid side effects.
- Have your child drink lots of noncaffeine liquids every day unless told to drink less liquid by your child's doctor.

What do I do if my child misses a dose?

- Call your child's doctor to find out what to do.

How do I store and/or throw out this drug?

- If you need to store this drug at home, talk with your child's doctor, nurse, or pharmacist about how to store it.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.

- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the

Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2025-03-19

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors.
All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Nelarabine - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering
Cancer Center