

Praziquantel

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Biltricide

Brand Names: Canada

Biltricide [DSC]

What is this drug used for?

- It is used to treat parasite worm infections.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has a worm infection of the eye.
- If your child has seizures or any other brain or nervous system problem.
- If your child has taken any of these drugs in the past 4 weeks: Carbamazepine, dexamethasone, efavirenz, phenobarbital, phenytoin, rifabutin, rifampin, or St. John's wort.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child avoid tasks or actions that call for alertness for 1 full day after your child takes this drug. These include things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized vehicles.
- If your child drinks grapefruit juice or eats grapefruit often, talk with your child's doctor.
- Some health problems have gotten worse during treatment with this drug. Sometimes, these may be life-threatening. If your child's signs do not get better or get worse, or if your child has any new signs during treatment, call your child's doctor right away.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Fever.
- Fast, slow, or abnormal heartbeat.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Feeling dizzy, tired, or weak.
- Headache.
- Stomach pain.
- Upset stomach.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- Give doses at least 4 hours apart but not more than 6 hours apart, unless told to do so by the doctor.
- Give this drug with a full glass of water.
- Give this drug with meals.
- The tablet may be broken if the doctor tells you to.

- Do not let your child chew this drug.
- Do not let your child hold the tablets or parts of the tablets in their mouth before swallowing. The bitter taste can lead to gagging or throwing up.
- If your child cannot swallow this drug whole, it may be crushed or dissolved and mixed with soft food or liquid. Give this drug within 1 hour of mixing.

What do I do if my child misses a dose?

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or

get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2024-02-02

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Praziquantel - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center

Last Updated

December 12, 2022

Learn about our [Health Information Policy](#).



Memorial Sloan Kettering
Cancer Center