PATIENT & CAREGIVER EDUCATION

Romiplostim

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Nplate

Brand Names: Canada

Nplate

What is this drug used for?

- It is used to raise platelet counts.
- It is used to treat immune thrombocytopenia (ITP).
- It is used to help with survival in some people who have had high amounts of radiation.
- It may be given to your child for other reasons. Talk with the doctor.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.

If your child is breast-feeding a baby:

- Be sure your child does not breast-feed a baby while taking this drug.

This drug may interact with other drugs or health problems.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.

- Use care to prevent your child from getting hurt and have your child avoid falls or crashes.
- Low platelet levels and a raised chance of bleeding may happen when this drug is stopped. If this drug is stopped, have your child's blood work checked for at least 2 weeks after your child stops taking it. If you have questions, talk with your child's doctor.
- This drug may raise the chance of blood clots. This includes some forms of blood clots that spread to the lungs or that cause strokes or heart attacks. Talk with the doctor.
- This drug is not for use in people with a certain bone marrow problem called myelodysplastic syndrome (MDS). If your child has MDS and takes this drug, MDS may get worse and turn into a type of blood cancer called acute myelogenous leukemia (AML). If MDS turns into AML, your child could die sooner from AML. If your child has MDS, talk with your child's doctor.
- If your child's weight changes, talk with the doctor. The dose of this drug may need to be changed.

If your child is pregnant:

- This drug may cause harm to the unborn baby if your child takes it during pregnancy. If your child is pregnant

or gets pregnant while taking this drug, call the doctor right away.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Any unexplained bruising or bleeding.
- Swelling in the arms or legs.
- Swelling of belly.
- Right-sided stomach pain.
- A burning, numbness, or tingling feeling that is not normal.
- Ear pain.
- Weakness on 1 side of the body, trouble speaking or

thinking, change in balance, drooping on one side of the face, or blurred eyesight.

- Call the doctor right away if your child has signs of a blood clot like chest pain or pressure; coughing up blood; shortness of breath; swelling, warmth, numbness, change of color, or pain in a leg or arm; or trouble speaking or swallowing.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Stomach pain or diarrhea.
- Signs of a common cold.
- Mouth or throat pain or irritation.
- Cough.
- Fever.
- Headache.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into the fatty part of the skin.

What do I do if my child misses a dose?

- Call your child's doctor to find out what to do.

How do I store and/or throw out this drug?

- If you need to store this drug at home, talk with your child's doctor, nurse, or pharmacist about how to store it.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do

so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health,

medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2025-04-14

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Romiplostim - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center

Memorial Sloan Kettering Cancer Center