

# Somatrogon

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

## Brand Names: US

Ngenla

## Brand Names: Canada

Ngenla

## What is this drug used for?

- It is used to help with growth and to treat growth hormone deficiency.
- It may be given to your child for other reasons. Talk with the doctor.

## What do I need to tell the doctor **BEFORE** my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has any of these health problems: Breathing problems; cancer; or illness shortly after open heart surgery, stomach surgery, or accidental injury.
- If your child's bones are no longer growing (closed epiphyses).
- If your child has eye problems caused by diabetes.
- If your child has Prader-Willi syndrome and is very overweight, has

trouble breathing, or has sleep apnea.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

## **What are some things I need to know or do while my child takes this drug?**

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child's blood work and other lab tests checked as you have been told by the doctor.
- High blood sugar has happened with drugs like this one. This includes diabetes that is new or worse.
- Check your child's blood sugar as you have been told by the doctor.
- This drug may affect certain lab tests. Tell all of your child's health care providers and lab workers that your child takes this drug.
- If your child has an abnormal curve in the spine (scoliosis), talk with your doctor. This drug could make it worse in children who are still growing.
- If your child has cancer or a tumor or has ever had cancer or a tumor, talk with your child's doctor. The chance of cancer or tumor growth is raised with this drug. The chance of new tumors may also be raised in some patients.
- Very bad and sometimes deadly lung or breathing problems have happened in some children with Prader-Willi syndrome. The chance may be higher in children who have sleep apnea, an infection in the lungs or airway, a block in the airway, and in children who are very overweight.

Call the doctor right away if your child has a fever of 100.4°F (38°C) or higher, cough, sore throat, shortness of breath, chest pain or discomfort, new or worse snoring, or breathing that is not normal while asleep.

## **If your child is pregnant or breast-feeding a baby:**

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

## **What are some side effects that I need to call my child's doctor about right away?**

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high blood sugar like confusion, feeling sleepy, unusual thirst or hunger, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain, or very bad upset stomach or throwing up.
- Signs of a weak adrenal gland like a severe upset stomach or throwing up, severe dizziness or passing out, muscle weakness, feeling very tired, mood changes, decreased appetite, or weight loss.
- Signs of low thyroid levels like constipation; not able to handle cold; memory problems; mood changes; or a burning, numbness, or tingling feeling that is not normal.
- Dizziness or passing out.
- Fast or abnormal heartbeat.

- Sweating a lot.
- Change in how your child acts.
- Change in eyesight.
- Change in skin color.
- A skin lump or growth.
- Change in color or size of a mole.
- Swelling.
- Muscle or joint pain.
- Burning, numbness, pain, or tingling in the hands, arms, legs, or feet.
- Ear pain or discharge.
- Fever.
- Feeling very tired or weak.
- Children who use this drug can rarely have a bone problem in the hip (slipped growth plate). Call the doctor right away if your child has hip or knee pain or a limp.
- Raised pressure in the brain has happened in children using drugs like this one. Call the doctor right away if your child has a headache; eyesight problems like blurred eyesight, seeing double, or loss of eyesight; pain behind the eye; pain when moving the eye; ringing in the ears; dizziness; or very upset stomach.

## **What are some other side effects of this drug?**

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Pain, redness, swelling, or other reaction where the injection was given.
- Nose or throat irritation.
- Headache.

- Cough.
- Throwing up.
- Signs of a common cold.
- Stomach pain.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

## **How is this drug best given?**

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into the fatty part of the skin in the upper arm, thigh, buttocks, or stomach area.
- If you will be giving your child the shot, your child's doctor or nurse will teach you how to give the shot.
- To make the injection feel better, take this drug out of the refrigerator and allow it to sit at room temperature for up to 30 minutes before use. Do not remove the cap or cover. Do not heat this drug.
- Move the site where you give the shot with each shot.
- If the dose is more than 1 injection, give the injections into 2 different places.
- Do not give into skin that is irritated, bruised, red, infected, hard, or scarred.
- Do not give into bony areas.
- Wash your hands before and after use.
- Do not shake the solution.
- Do not use if the solution is cloudy, leaking, or has particles.

- This drug is colorless to a faint yellow. Do not use if the solution changes color.
- This product may make a clicking sound as you prepare the dose. Do not prepare the dose by counting the clicks. Doing so could lead to using the wrong dose.
- Attach new needle before each dose.
- Remove all pen needle covers before injecting a dose (there may be 2). If you are not sure what type of pen needle you have or how to use it, talk with the doctor.
- Do not share pen or cartridge devices with another person even if the needle has been changed. Sharing these devices may pass infections from one person to another. This includes infections you may not know your child has.
- Take off the needle after each shot. Do not store this device with the needle on it.
- Put the cap back on after you are done using your dose.
- Throw away needles in a needle/sharp disposal box. Do not reuse needles or other items. When the box is full, follow all local rules for getting rid of it. Talk with a doctor or pharmacist if you have any questions.
- Do not use this drug if it has been dropped or if it is broken.

## **What do I do if my child misses a dose?**

- Give a missed dose as soon as you think about it.
- If your child misses a dose by more than 3 days, skip the missed dose. Give the next dose on your child's normal day.
- Do not give 2 doses within 3 days of each other.

## How do I store and/or throw out this drug?

- Store unopened and opened pens in a refrigerator. Do not freeze.
- Store in the outer carton to protect from light.
- Protect from heat.
- Do not use if it has been frozen.
- After opening, throw away any part not used after 28 days.
- Keep out of direct sunlight.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

## General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

## Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

## Last Reviewed Date

2023-07-24

## Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.



If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

Somatrogon - Last updated on January 10, 2023

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering  
Cancer Center