

PATIENT & CAREGIVER EDUCATION

Somatropin

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Genotropin; Genotropin MiniQuick; Humatrope; Norditropin FlexPro; Nutropin AQ NuSpin 10; Nutropin AQ NuSpin 20 [DSC]; Nutropin AQ NuSpin 5 [DSC]; Omnitrope; Saizen [DSC]; Saizenprep [DSC]; Serostim; Zomacton; Zomacton (for Zoma-Jet 10) [DSC]; Zorbtive [DSC]

Brand Names: Canada

Genotropin GoQuick; Genotropin MiniQuick; Humatrope [DSC]; Norditropin FlexPro; Norditropin NordiFlex Pen [DSC]; Nutropin AQ NuSpin 10 [DSC]; Nutropin AQ NuSpin 20 [DSC]; Nutropin AQ NuSpin 5 [DSC]; Omnitrope; Saizen; Serostim

What is this drug used for?

- It is used to help with growth and to treat growth hormone deficiency.
- It may be given to your child for other reasons. Talk with the doctor.

What do I need to tell the doctor **BEFORE** my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has any of these health problems: Breathing problems like sleep apnea; cancer or other tumors like a brain tumor; diabetic eye disease; or illness shortly after open heart surgery, stomach surgery, or accidental injury.
- If your child has Prader-Willi syndrome and is very overweight, has trouble breathing, or has sleep apnea.
- If your child's bones are no longer growing (closed epiphyses).

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins)

and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- High blood sugar has happened with this drug. This includes diabetes that is new or worse.
- Check your child's blood sugar as you have been told by the doctor.
- Have your child's blood work checked and eye exams as you have been told by your child's doctor.
- This drug may affect certain lab tests. Tell all of your child's health care providers and lab workers that your child takes this drug.
- If your child has an abnormal curve in the spine (scoliosis), talk with your doctor. This drug could make it worse in children who are still growing.
- If your child has Turner syndrome, talk with the doctor.

The chance of ear infections, high blood pressure, and very bad blood vessel problems like stroke and bleeding in the brain may be raised.

- If your child has cancer or a tumor or has ever had cancer or a tumor, talk with your child's doctor. The chance of cancer or tumor growth is raised with this drug. The chance of new tumors may also be raised in some patients.
- Some products have benzyl alcohol. If possible, avoid products with benzyl alcohol in newborns or infants. Serious side effects can happen in these children with some doses of benzyl alcohol, including if given with other drugs that have benzyl alcohol. Talk with the doctor to see if this product has benzyl alcohol in it.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side

effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high blood sugar like confusion, feeling sleepy, unusual thirst or hunger, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain, or very bad upset stomach or throwing up.
- Signs of a weak adrenal gland like a severe upset stomach or throwing up, severe dizziness or passing out, muscle weakness, feeling very tired, mood changes, decreased appetite, or weight loss.
- Signs of low thyroid levels like constipation; not able to handle cold; memory problems; mood changes; or a burning, numbness, or tingling feeling that is not normal.
- Signs of a urinary tract infection (UTI) like blood in the urine, burning or pain when passing urine, feeling the

need to pass urine often or right away, fever, lower stomach pain, or pelvic pain.

- Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight.
- Shortness of breath, a big weight gain, or swelling in the arms or legs.
- Chest pain or pressure or a fast heartbeat.
- Sweating a lot.
- Depression or other mood changes.
- Change in how you act.
- Change in skin color.
- Burning, numbness, pain, or tingling in the hands, arms, wrists, legs, or feet.
- Bone pain.
- Change in color or size of a mole.
- Redness or swelling where the injection is given.
- Skin breakdown where this drug is used.
- Ear pain.
- Children who use this drug may rarely have a bone

problem in the hip (slipped growth plate). This can also lead to further bone injury. Call the doctor right away if your child has hip or knee pain or a limp.

- Raised pressure in the head has rarely happened with this drug. The risk may be greater in patients with Turner syndrome or Prader-Willi syndrome. Signs most often happen within the first 8 weeks of starting this drug. Call your child's doctor right away if your child has a change in eyesight, a severe headache, upset stomach, or throwing up.
- Very bad and sometimes deadly lung or breathing problems have happened in some children with Prader-Willi syndrome. The chance may be higher in children who have sleep apnea, an infection in the lungs or airway, a block in the airway, and in children who are very overweight. Call the doctor right away if your child has a fever of 100.4°F (38°C) or higher, cough, sore throat, shortness of breath, chest pain or discomfort, new or worse snoring, or breathing that is not normal while asleep.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do

not go away:

- Headache.
- Feeling tired or weak.
- Back, muscle, or joint pain.
- Muscle stiffness.
- Trouble sleeping.
- Diarrhea, stomach pain, upset stomach, or throwing up.
- Gas.
- Irritation where the shot is given.
- Signs of a common cold.
- Flu-like signs.
- Hair loss.
- Enlarged breasts.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

All products:

- It is given as a shot into the fatty part of the skin.
- If you will be giving your child the shot, your child's doctor or nurse will teach you how to give the shot.
- If stored in a refrigerator, let this drug come to room temperature before using it. Be sure you know how long to leave it at room temperature before using. Do not heat this drug.
- Move the site where you give the shot with each shot.
- Do not give into skin that is irritated, bruised, red, infected, hard, or scarred.
- Do not give into skin near the belly button or waistline.
- Wash your hands before and after use.
- Do not shake the solution.
- Do not use if the solution is cloudy, leaking, or has particles.
- Do not use if solution changes color.
- Throw away needles in a needle/sharp disposal box. Do not reuse needles or other items. When the box is full,

follow all local rules for getting rid of it. Talk with a doctor or pharmacist if you have any questions.

Prefilled syringes or pens:

- This product may make a clicking sound as you prepare the dose. Do not prepare the dose by counting the clicks. Doing so could lead to using the wrong dose.
- Remove all pen needle covers before injecting a dose (there may be 2). If you are not sure what type of pen needle you have or how to use it, talk with the doctor.
- Do not share pen or cartridge devices with another person even if the needle has been changed. Sharing these devices may pass infections from one person to another. This includes infections you may not know your child has.

Cartridge injection devices (pens):

- Not all injection devices (pens) are for use with all strengths of this drug. Be sure you have the right pen to use with the cartridge for your child's dose. Using the wrong pen may give your child the wrong dose.

What do I do if my child misses a dose?

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip

the missed dose and go back to your child's normal time.

- Do not give 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Some brands of this drug need to be stored in a refrigerator. Some brands of this drug need to be stored at room temperature. If you have questions about how to store this drug, talk with your pharmacist.
- If stored in the refrigerator, do not store right next to the cooling element.
- Protect from light.
- Be sure you know how long you can store this drug before you need to throw it away.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.

- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any

risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2025-07-30

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors.
All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Somatropin - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering
Cancer Center