

Sparsentan

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Filspari

Warning

- Severe liver problems may happen with this drug. This drug may not be right for patients who already have raised liver enzymes or liver problems. Tell the doctor if your child has liver problems. Your child will need to have blood work checked before starting this drug and while taking it. If you have questions, talk with the doctor.
- Call the doctor right away if your child has signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach, stomach pain, light-colored stools, throwing up, or yellow skin or eyes.

- This drug may cause harm to the unborn baby if your child takes it during pregnancy.
- Do not let your child take this drug during pregnancy. A pregnancy test will be done to show that your child is NOT pregnant before starting this drug.
- If your child may become pregnant, birth control must be used before starting this drug, while taking it, and for at least 2 weeks after the last dose. If your child becomes pregnant, call the doctor right away.
- If your child misses a period, has unprotected sex, or thinks that birth control has not worked, call the doctor right away.

What is this drug used for?

- It is used to lower protein in the urine in certain patients with focal segmental glomerulosclerosis (FSGS).
- If your child has been given this drug for some other reason, talk with the doctor about the benefits and risks. Talk with the doctor if you have questions or concerns about giving this drug to your child.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child takes any drugs (prescription or OTC, natural products, vitamins) that must not be taken with this drug, like certain drugs that are used for HIV, infections, depression, high blood pressure, and others. There are many drugs that must not be taken with this drug. The doctor or pharmacist can tell you if your child is taking a drug that must not be taken with this drug.
- If your child is breast-feeding a baby. Be sure your child does not breast-feed while taking this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- To lower the chance of feeling dizzy or passing out, have your child rise slowly if your child has been sitting or lying down. Have your child be careful going up and down stairs.
- Have blood work and blood pressure checked as you have been told by the doctor.
- If your child drinks grapefruit juice or eats grapefruit often, talk with your child's doctor.
- If your child is taking a salt substitute that has potassium in it, a potassium-sparing diuretic, or a potassium product, talk with your child's doctor.
- If your child started taking this drug before puberty, talk with the doctor. Call the doctor right away if breast buds or pubic hair begin to develop or if your child's period starts.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high potassium levels like a heartbeat that does not feel normal; feeling confused; feeling weak, lightheaded, or dizzy; feeling like passing out; numbness or tingling; or shortness of breath.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Severe dizziness or passing out.
- Shortness of breath, a big weight gain, or swelling in the arms or legs.
- Feeling very tired or weak.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Dizziness.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- Give before the morning or the evening meal. Always give it before the same meal every day.
- Have your child swallow whole with a drink of water.
- If your child cannot swallow this drug whole, it may be crushed and mixed with $\frac{1}{2}$ cup (120 mL) of water. The crushed tablet will not dissolve. Have your child drink right away after mixing. Do not store for use at a later

time. If any of this drug is left after drinking, add a little more water to the cup, swirl, and have your child swallow.

- Have your child drink lots of noncaffeine liquids unless told to drink less liquid by your child's doctor.

What do I do if my child misses a dose?

- Skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store in the original container at room temperature.
- Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination

and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2026-04-28

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors.

All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Sparsentan - Last updated on April 30, 2026

All rights owned and reserved by Memorial Sloan Kettering Cancer Center

Last Updated

April 30, 2026

Learn about our [Health Information Policy](#).



Memorial Sloan Kettering
Cancer Center