

Triptorelin

This information from UpToDate® Lexidrug™ explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Trelstar Mixject; Triptodur

Brand Names: Canada

Decapeptyl; Trelstar

What is this drug used for?

- It is used to delay puberty in children who are maturing too early.
- If your child has been given this drug for some other reason, talk with the doctor about the benefits and risks. Talk with the doctor if you have questions or concerns about giving this drug to your child.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.

If your child is or may be pregnant:

- If your child is pregnant or may be pregnant. Do not give this drug to your child during pregnancy.

This drug may interact with other drugs or health problems.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- This drug may affect certain lab tests. Tell all of your child's health care providers and lab workers that your child takes this drug.
- This drug may affect growth in children and teens in some cases. They may need regular growth checks. Talk with the doctor.

If your child is pregnant:

- This drug may cause harm to the unborn baby if your child takes it during pregnancy. If your child is pregnant or gets pregnant while taking this drug, call the doctor right away.

If your child is breast-feeding a baby:

- Tell the doctor if your child is breast-feeding a baby. You will need to talk about any risks to the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs

or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal.
- Seizures.
- A fast heartbeat.
- Dizziness or passing out.
- Sweating a lot.
- Diarrhea.
- Stomach pain.
- Stomach cramps.
- Upset stomach or throwing up.
- Behavior and mood changes have happened with the use of drugs like this one in children. This includes acting aggressive, crying, depression, emotional ups and downs, restlessness, feeling angry and irritable, and suicidal thoughts or actions. Call your child's doctor right away if your child has any new or worse behavior or mood changes.
- Raised pressure in the brain has happened with this drug. Call your child's doctor right away if your child has a headache; dizziness; upset stomach; ringing in the ears; eye pain; or eyesight problems like blurred eyesight, seeing double, or loss of eyesight.
- Severe skin reactions may happen with this drug. These include Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and other serious reactions. Sometimes, body organs may also be affected. These reactions can be deadly. Get medical help right away if your child has signs like red, swollen, blistered, or peeling skin; red or irritated eyes; sores in the mouth,

throat, nose, eyes, genitals, or any areas of skin; fever; chills; body aches; shortness of breath; or swollen glands.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Nose or throat irritation.
- Signs of a common cold.
- Headache.
- Hot flashes.
- Irritation where the shot is given.
- If your child is taking this drug to delay puberty, you may see short-term body odor or signs of puberty like vaginal bleeding, enlarged breasts, or more pubic hair during the first month of care. If these last or are very bad, call the doctor.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into a muscle.

What do I do if my child misses a dose?

- Call your child's doctor to find out what to do.

How do I store and/or throw out this drug?

- If you need to store this drug at home, talk with your child's doctor, nurse, or pharmacist about how to store it.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2025-10-21

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Triptorelin - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center

Last Updated

December 12, 2022

Learn about our [Health Information Policy](#).



Memorial Sloan Kettering
Cancer Center