



PATIENT & CAREGIVER EDUCATION

Metreleptin

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Myalept

Warning

- Some people who use this drug make antibodies in their blood. These may lower how well a hormone (leptin) in the body works. They may also lower how well this drug works. Side effects may also happen. These could be infection, problems with blood sugar like diabetes, and raised fats in the blood. Talk with the doctor.
- A certain type of lymphoma has happened in people with this health problem. This has happened in people both treated and not treated with this drug. Tell the doctor if your child has ever had any immune system or bone marrow problems. Talk with the doctor.

What is this drug used for?

- It is used to treat problems caused by not having enough of a certain hormone (leptin) in the body. It is only for use in people with congenital or acquired generalized lipodystrophy.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.

- If your child's health problem is not caused by not having enough leptin in the body.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for your child to take this drug with all of his/her drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child's blood work checked often. Talk with your child's doctor.
- Have your child follow the diet plan your child's doctor told you about.
- Do not stop giving this drug to your child all of a sudden without calling the doctor. Your child may have a greater risk of side effects. If your child needs to stop this drug, you will want to slowly stop it as told by the doctor.
- If your child has had an autoimmune disease your child's signs may get worse. Talk with the doctor.
- This drug may lower your child's blood sugar levels. The chance may be raised if your child also takes insulin or other drugs that lower blood sugar. Talk with the doctor.
- Have your child's blood sugar checked as you have been told by your child's doctor.
- Some fluids used to mix this drug have benzyl alcohol. Do not give a product that has benzyl alcohol in it to a newborn or infant. It may cause very bad and sometimes deadly side effects. Talk with the doctor to see which fluids have benzyl alcohol.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is

breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal.
- Signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- Dizziness or passing out.
- A fast heartbeat.
- A burning, numbness, or tingling feeling that is not normal.
- Pelvic pain or pressure.
- Vaginal bleeding that is not normal.
- Call your child's doctor right away if your child has a swollen gland, night sweats, shortness of breath, or weight loss without trying.
- Low blood sugar can happen. The chance may be raised when this drug is used with other drugs for diabetes. Signs may be dizziness, headache, feeling sleepy or weak, shaking, fast heartbeat, confusion, hunger, or sweating. Call your doctor right away if you have any of these signs. Follow what you have been told to do for low blood sugar. This may include taking glucose tablets, liquid glucose, or some fruit juices.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Headache.
- Stomach pain or diarrhea.
- Weight loss.
- Joint pain.
- Ear pain.
- Feeling tired or weak.
- Upset stomach.
- Signs of a common cold.
- Back pain.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into the fatty part of the skin on the top of the thigh, belly area, or upper arm.
- If you will be giving your child the shot, your child's doctor or nurse will teach you how to give the shot.
- Wash your hands before and after use.
- This drug needs to be mixed before use. Follow how to mix as you were told by the doctor.
- A few fluids can be used to mix this drug. Be sure you know which fluid to use.

Talk with your doctor or pharmacist if you are not sure.

- Let this drug come to room temperature before mixing. Do not heat this drug.
- Do not shake.
- Move site where you give the shot each time.
- Give this drug with or without food.
- Give this drug at the same time of day.
- Do not mix with any other liquid drugs.
- Do not mix this drug in the same syringe with insulin.
- If your child is also using insulin, you may inject this drug and the insulin in the same area of the body but not right next to each other.
- Do not use if the solution is cloudy, leaking, or has particles.
- Do not use if solution changes color.
- Throw away needles in a needle/sharp disposal box. Do not reuse needles or other items. When the box is full, follow all local rules for getting rid of it. Talk with a doctor or pharmacist if you have any questions.

What do I do if my child misses a dose?

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store unopened vials in a refrigerator. Do not freeze.
- Store in the original container to protect from light.
- After mixing, be sure you know how long the product is good for and how to store it. Ask your doctor or pharmacist if you are not sure.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.

- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of using this medicine.

Last Reviewed Date

2020-01-30

Copyright

© 2020 Wolters Kluwer Clinical Drug Information, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your healthcare team directly. If you're a patient at MSK and you need to reach a provider after 5:00 PM, during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Metreleptin - Last updated on February 11, 2020

©2020 Memorial Sloan Kettering Cancer Center