



## PATIENT & CAREGIVER EDUCATION

# Minocycline

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

### **Brand Names: US**

CoreMino; Minocin; Minolira; Solodyn; Ximino

### **What is this drug used for?**

- It is used to treat or prevent bacterial infections.
- It is used to treat pimples (acne).
- It may be given to your child for other reasons. Talk with the doctor.

### **What do I need to tell the doctor BEFORE my child takes this drug?**

#### **All products:**

- If your child has an allergy to this drug or any part of this drug.
- If your child is allergic to any drugs like this one or any other drugs, foods, or other substances. Tell the doctor about the allergy and what signs your child had, like rash; hives;

itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs.

- If your child is taking any of these drugs: Acitretin, isotretinoin, or a penicillin.

### **All extended-release products:**

#### **If your child is or may be sexually active:**

- If your child may become pregnant or father a child.

#### **If your child is pregnant or breast-feeding a baby:**

- Do not give this drug to your child if she is pregnant.
- Talk with the doctor if your child is breast-feeding a baby or plans to breast-feed a baby.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for your child to take this drug with all of his/her drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

### **What are some things I need to know or do while my child takes this drug?**

#### **All products:**

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Your child may get sunburned more easily. Avoid lots of sun, sunlamps, and tanning beds. Use sunscreen and dress your child in clothing and eyewear that protects him/her from the sun.
- Have your child avoid tasks or actions that call for alertness until you see how this drug affects your child. These are things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized vehicles.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- This drug may affect certain lab tests. Tell all of your child's health care providers and lab workers that your child takes this drug.
- Do not give to your child longer than you have been told. A second infection may happen.
- If your child is allergic to sulfites, talk with your child's doctor. Some products have sulfites in them.
- A very bad and sometimes deadly reaction has happened with this drug. Most of the time, this reaction has signs like fever, rash, or swollen glands with problems in body organs like the liver, kidney, blood, heart, muscles and joints, or lungs. Talk with the doctor.

- A very bad skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if your child has signs like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in the mouth, throat, nose, or eyes.
- This drug may cause a change in tooth color to yellow-gray-brown in children younger than 8 years old. If this change of tooth color happens, it will not go away. Talk with the doctor.

**If your child is or may be sexually active:**

- Birth control pills and other hormone-based birth control may not work as well to prevent pregnancy. Be sure your child uses some other kind of birth control also, like a condom, when taking this drug.

**If your child is pregnant:**

- This drug may cause harm to the unborn baby if your child takes it during pregnancy. If your child is pregnant or gets pregnant while taking this drug, call the doctor right away.

**All extended-release products:**

- This drug is not approved for use in children younger than 12 years old. Talk with the doctor.

- Do not give to a child younger than 8 years of age.

### **All other products:**

- Most of the time, this drug is not for use in children younger than 8 years old. However, there may be times when these children may need to take this drug. Talk with the doctor.

### **What are some side effects that I need to call my child's doctor about right away?**

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain, or very bad upset stomach or throwing up.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Signs of lupus like a rash on the cheeks or other body parts, sunburn easy, muscle or joint pain, chest pain or shortness

of breath, or swelling in the arms or legs.

- Swollen gland.
- Fever, chills, or sore throat; any unexplained bruising or bleeding; or feeling very tired or weak.
- Change in hearing.
- Joint pain or swelling.
- Muscle pain or weakness.
- Ringing in ears.
- Seizures.
- Shortness of breath.
- Chest pain.
- A heartbeat that does not feel normal.
- Swelling.
- Change in color of nails, skin, eyes, scars, teeth, or gums to a darker color.
- Mouth irritation or mouth sores.
- A burning, numbness, or tingling feeling that is not normal.
- Redness or white patches in mouth or throat.
- Rectal irritation.
- Genital irritation.
- Vaginal itching or discharge.
- It is common to have diarrhea when taking antibiotics.

Rarely, a severe form of diarrhea called C diff-associated diarrhea (CDAD) may happen. Sometimes, this has led to a deadly bowel problem (colitis). CDAD may happen while your child is taking an antibiotic or within a few months after he/she stops taking it. Call your child's doctor right away if your child has stomach pain or cramps, very loose or watery stools, or bloody stools. Do not try to treat diarrhea without first checking with the doctor.

- Very bad and sometimes deadly liver problems have happened with this drug. Call your child's doctor right away if your child has signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Raised pressure in the brain has happened with this drug. Most of the time, this will go back to normal after this drug is stopped. Sometimes, loss of eyesight may happen and may not go away even after this drug is stopped. Call the doctor right away if your child has a headache or eyesight problems like blurred eyesight, seeing double, or loss of eyesight.
- Cases of thyroid cancer have been reported with long-term use of this drug. Call your child's doctor right away if your child has lasting hoarseness, a neck mass, or trouble breathing or swallowing.

### **What are some other side effects of this drug?**

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's

doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

### **All products:**

- Diarrhea, upset stomach, or throwing up.
- Not hungry.
- Feeling dizzy, sleepy, tired, or weak.

### **Injection:**

- Irritation where the shot is given.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

### **How is this drug best given?**

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

### **All oral products:**

- Keep using this drug as you have been told by your child's doctor or other health care provider, even if your child's signs get better.
- Give this drug at the same time of day.
- Give this drug with or without food. Give with food if it causes an upset stomach.

- Do not give products that have iron in them or products like antacids that have aluminum, calcium, or magnesium in them at the same time as this drug. Talk with the doctor or pharmacist.
- Have your child drink lots of noncaffeine liquids every day unless told to drink less liquid by your child's doctor.

### **Tablets and capsules:**

- Have your child swallow whole with a full glass of water.

### **All extended-release products:**

- Have your child swallow whole. Do not let your child chew or crush.
- Give this drug with a full glass of water.

### **Extended-release tablets:**

- Some products may be broken in half. If you are not sure if you can break this product in half, talk with the doctor.

### **Liquid (suspension):**

- Shake well before use.
- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.

### **Injection:**

- It is given as an infusion into a vein over a period of time.

## **What do I do if my child misses a dose?**

### **All oral products:**

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

### **Injection:**

- Call your child's doctor to find out what to do.

## **How do I store and/or throw out this drug?**

### **All oral products:**

- Store at room temperature.
- Store in a dry place. Do not store in a bathroom.
- Protect from heat and light.
- Get rid of this drug when your child no longer needs it or if the drug is outdated.

### **Injection:**

- If you need to store this drug at home, talk with your child's doctor, nurse, or pharmacist about how to store it.

### **All products:**

- Keep all drugs in a safe place. Keep all drugs out of the reach

of children and pets.

- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

## **General drug facts**

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Keep a list of all your child's drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your child's doctor.
- Talk with your child's doctor before giving your child any new drug, including prescription or OTC, natural products, or vitamins.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

## **Consumer Information Use and Disclaimer**

This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of using this medicine.

### **Last Reviewed Date**

2019-06-11

### **Copyright**

© 2019 Wolters Kluwer Clinical Drug Information, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your healthcare team directly. If you're a patient at MSK and you need to reach a provider after 5:00 PM, during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

---

Minocycline - Generated on July 18, 2019

©2019 Memorial Sloan Kettering Cancer Center