



PATIENT & CAREGIVER EDUCATION

Molindone

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Warning

- There is a higher chance of death in older adults who take this drug for mental problems caused by dementia. Most of the deaths were linked to heart disease or infection. This drug is not approved to treat mental problems caused by dementia.

What is this drug used for?

- It is used to treat schizophrenia.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take this drug?

For all patients taking this drug:

- If you have an allergy to molindone or any other part of this drug.
- If you are allergic to any drugs like this one, any other drugs, foods, or other substances. Tell your doctor about the allergy and what signs you had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs.
- If you are very sleepy.
- If you have recently drunk a lot of alcohol or taken a big amount of drugs that may slow your actions like phenobarbital or some pain drugs like oxycodone.

Children:

- If your child is younger than 12 years of age. Do not give this drug to a child younger than 12 years of age.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.
- Do not stop taking this drug all of a sudden without calling your doctor. You may have a greater risk of side effects. If you need to stop this drug, you will want to slowly stop it as ordered by your doctor.
- This drug may prevent other drugs taken by mouth from getting into the body. If you take other drugs by mouth, you may need to take them at some other time than this drug. Talk with your doctor.
- Low white blood cell counts have happened with drugs like this one. This may lead to a higher chance of getting an infection. Deadly infections have rarely happened. Tell your doctor if you have ever had a low white blood cell count. Call your doctor right away if you have signs of infection like fever, chills, or sore throat. Talk with your doctor.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- Talk with your doctor before you drink alcohol or use other drugs and natural products that slow your actions.

- Dizziness, sleepiness, and feeling less stable may happen with this drug. These may lead to falling. Broken bones or other health problems can happen from falling. Talk with the doctor.
- Older adults with dementia taking drugs like this one have had a higher number of strokes. Sometimes these strokes have been deadly. This drug is not approved to treat mental problems caused by dementia. Talk with your doctor.
- If you are 65 or older, use this drug with care. You could have more side effects.
- Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.
- Taking this drug in the third trimester of pregnancy may lead to muscle movements that cannot be controlled and withdrawal in the newborn. Talk with the doctor.
- Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Shakiness, trouble moving around, or stiffness.
- Seizures.
- Nipple discharge.
- For women, no period.

- Enlarged breasts.
- Not able to get or keep an erection.
- More interest in sex.
- Not able to sit still.
- Mood changes.
- Restlessness.
- Trouble swallowing.
- A fast heartbeat.
- Trouble passing urine.
- Some people who take this drug may get a very bad muscle problem called tardive dyskinesia. This muscle problem may not go away even if this drug is stopped. Sometimes, signs may lessen or go away over time after this drug is stopped. The risk of tardive dyskinesia may be greater in people with diabetes and in older adults, especially older women. The risk is also greater the longer you take this drug or with higher doses. Muscle problems may also occur after short-term use with low doses. Call your doctor right away if you have trouble controlling body movements or if you have muscle problems with your tongue, face, mouth, or jaw like tongue sticking out, puffing cheeks, mouth puckering, or chewing.
- A very bad and sometimes deadly health problem called neuroleptic malignant syndrome (NMS) may happen. Call your doctor right away if you have any fever, muscle cramps or stiffness, dizziness, very bad headache, confusion, change in thinking, fast heartbeat, heartbeat that does not feel normal, or are sweating a lot.
- Call your doctor right away if you have a painful erection (hard penis) or an erection that lasts for longer than 4 hours. This may happen even when you are not having sex. If this is not treated right away, it may lead to lasting sex problems and you may not be able to have sex.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Feeling sleepy.
- Blurred eyesight.
- Upset stomach.
- Dry mouth.
- More saliva.
- Constipation.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- To gain the most benefit, do not miss doses.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature.
- Store in a dry place. Do not store in a bathroom.
- Protect from light.
- Keep lid tightly closed.

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Keep a list of all your drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your doctor.
- Talk with the doctor before starting any new drug, including prescription or OTC, natural products, or vitamins.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of using this

medicine.

Last Reviewed Date

2017-08-16

Copyright

© 2019 Wolters Kluwer Clinical Drug Information, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your healthcare team directly. If you're a patient at MSK and you need to reach a provider after 5:00 PM, during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Molindone - Generated on June 26, 2019

©2019 Memorial Sloan Kettering Cancer Center