

Outpatient Bispecific Antibody Emergency Guide

| | Major Emergencies: Call 911 | Non-Emergencies: Call your doctor's office |
|----------------------------------|--|--|
| Alertness | <ul style="list-style-type: none"> • Will not wake up • Suddenly cannot move arms or legs • Seizure (uncontrolled shaking or stiffening of body) • Sudden change in vision | <ul style="list-style-type: none"> • Dizziness or lightheadedness • Tremors or shakiness • Changes in mood • Change in energy level • New confusion or more confusion than usual • Trouble finding words • Change in level of alertness • Numbness or tingling • Unable to get around • Trouble swallowing |
| Breathing | <ul style="list-style-type: none"> • Not breathing • Trouble breathing | <ul style="list-style-type: none"> • Wheezing (whistling or squeaking sound when breathing) • Gets winded (short of breath) more easily with very little activity • New cough or cough that will not go away • Coughing up blood or green or yellow mucus (a jelly-like substance) • Trouble breathing when lying down |
| Fever/chills or infection | | <ul style="list-style-type: none"> • Fever of 100.4 °F (38 °C) or higher, when taking temperature by mouth • Shaking chills (temperature may be normal) • Cold symptoms (runny nose, watery eyes, sneezing, coughing) • Pain or burning while urinating (peeing) |

