Memorial Sloan Kettering Cancer Center

PATIENT & CAREGIVER EDUCATION

Palliative Radiation Therapy to Your Spine

This information explains what to expect during your simulation for palliative radiation therapy to your spine. It also explains what to expect during your treatments and side effects you may have.

Palliative radiation therapy is radiation therapy that's meant to help ease some types of cancer symptoms. Your radiation therapy care team will talk with you about the goals of your palliative radiation therapy.

Simulation

Simulation (also called a mapping or planning session) is a procedure to plan your radiation treatments. It's done to map your treatment site, make sure you get the right dose (amount) of radiation, and limit the amount of radiation that gets to your nearby tissues.

During your simulation, we may make a mold of your upper body or pelvis (see Figure 1). You'll use the mold during your simulation and all your treatments to help you stay still

and make sure you're in the same position every time. Making the mold will take about 15 minutes.




Figure 1. Examples of radiation therapy molds

Take devices off your skin

You may wear certain devices on your skin. Before your simulation or treatment, some device makers recommend you take off your:

- Continuous glucose monitor (CGM)
- Insulin pump

If you use one of these, ask your radiation oncologist if you need to take it off. If you do, make sure to bring an extra device to put on after your simulation or treatment.

You may not be sure how to manage your glucose (blood sugar) while your device is off. If so, before your

appointment, talk with the healthcare provider who manages your diabetes care.

What to expect during each radiation treatment

A staff member will bring you to the treatment room. There, the radiation therapist will greet you and help you onto the treatment table (see Figure 2). You'll be positioned exactly how you were during your simulation. Then, you'll get your treatment.




Figure 2. Examples of radiation treatment machines

You'll be in the treatment room for 15 to 90 minutes, depending on your treatment plan. Most of this time will be spent making sure you're in the right position. The radiation therapist will walk in and out of the room, but they'll be able to see and hear you the whole time.

Side effects of palliative radiation therapy to your spine

You'll probably have some side effects during and after your treatments. Common side effects are listed below.

If you have any side effects, be sure to tell a member of your radiation therapy team. We'll help you manage them.

- **Pain flare and swelling around the treatment site:** Pain flare is a temporary increase in pain. During your treatment, the treatment area may become swollen and painful. We may prescribe medicine to help with this.
- **Nausea or vomiting**: Nausea is when you feel like you might vomit (throw up). You may have nausea, vomiting, or both during your treatment. We may prescribe an anti-nausea medicine to help prevent this. You'll take the medicine 30 minutes before each of your treatments.
- **Esophagitis:** Esophagitis (ee-SAH-fuh-JY-tis) is when your esophagus (food pipe) is irritated or swollen. It can make it difficult or painful to swallow. If your esophagus is in the treatment area, you may have esophagitis starting about 1 week after your treatment ends. Esophagitis can last for up to 1 week. We'll talk with you about ways to manage it.

- **Bowel changes:** During your treatment, your bowel movements (poop) may be softer or happen more often than usual.
- **Skin reactions:** Your skin may become pink or tanned about 2 to 3 weeks into your treatment. As you continue treatment, it may become bright red or very dark. It might also feel dry and itchy and look flaky. You can use lotions meant for dry skin (such as Eucerin®) to help keep it moisturized. Protecting your skin from the sun by wearing sunscreen will also help. These skin reactions will gradually get better about 3 or 4 weeks after your treatment ends.
- **Fatigue:** Fatigue is a feeling of being tired or weak, not wanting to do things, not being able to concentrate, or feeling slowed down. Fatigue can range from mild to severe. You may develop fatigue during or after your treatment, and it may last for weeks to months after your treatment ends. Other treatments you're getting can affect how long fatigue lasts.

Simulation date:

_____

Treatment start date:

_____

Medicine instructions:

_____

For more resources, visit www.mskcc.org/pe to search our virtual library.

Palliative Radiation Therapy to Your Spine - Last updated on February 27, 2026

All rights owned and reserved by Memorial Sloan Kettering Cancer Center