



## PATIENT & CAREGIVER EDUCATION

# About Palliative Whole Brain Radiation Therapy

This information explains what to expect during your simulation for palliative whole brain radiation therapy. It also explains what to expect during your treatments and side effects you may have.

Palliative radiation therapy is radiation therapy that's meant to help ease some types of cancer symptoms. Your radiation therapy team will talk with you about the goals of your palliative radiation therapy.

## Simulation

### Take devices off your skin

You may wear certain devices on your skin. Before your simulation or treatment, some device makers recommend you take off your:

- Continuous glucose monitor (CGM)
- Insulin pump

If you use one of these, ask your radiation oncologist if you

need to take it off. If you do, make sure to bring an extra device to put on after your simulation or treatment.

You may not be sure how to manage your glucose (blood sugar) while your device is off. If so, before your appointment, talk with the healthcare provider who manages your diabetes care.

### **During your simulation**

Simulation (also called a mapping or planning session) is a procedure to plan your radiation treatments. It's done to map your treatment site, make sure you get the right dose (amount) of radiation, and limit the amount of radiation that gets to your nearby tissues.

If you wear a head covering (such as a wig, turban, or cap), you may need to take it off before your simulation. If you use a hearing aid, ask your radiation therapist if you need to take it out.

During your simulation, we may make a plastic mesh mold of your face and head (see Figure 1). You'll use the mold during your simulation and all your treatments to help you stay still and make sure your head is in the same position every time. You'll be able to see and hear with it on. Making




Figure 1. An example of a mold used for whole brain radiation therapy

the mold will take about 15 minutes.

## What to expect during each radiation treatment

A staff member will bring you to the radiation treatment room. There, the radiation therapist will greet you and help you onto the treatment table (see Figure 2). You'll be positioned exactly how you were during your simulation. Then, you'll get your treatment. During your treatment, the machine will move in a circle around you.




Figure 2. An example of a radiation therapy machine

---

You'll be in the treatment room for 15 to 90 minutes,

depending on your treatment plan. Most of this time will be spent making sure you're in the right position. The radiation therapist will walk in and out of the room, but they'll be able to see and hear you the whole time.

## **Side effects of palliative whole brain radiation therapy**

You'll probably have some side effects during and after your treatments. Common side effects are listed below.

If you have any side effects, be sure to tell a member of your radiation therapy team. We'll help you manage them.

- **Hair loss:** You'll probably lose some of the hair on your head and eyebrows about 2 to 3 weeks into your treatment. Your hair should grow back 3 to 6 months after your treatment ends.
- **Skin reactions:** Your skin may become pink or tanned about 2 to 3 weeks into your treatment. As you continue treatment, it may become bright red or very dark. It might also feel dry and itchy and look flaky. You can use lotions meant for dry skin (such as Eucerin®) to help keep it moisturized. Protecting it from the sun by wearing sunscreen will also help. These skin reactions will start to gradually get better about 3 or 4 weeks after your treatment ends.

- **Fatigue:** Fatigue is a feeling of being tired or weak, not wanting to do things, not being able to concentrate, or feeling slowed down. Fatigue can range from mild to severe. You may develop fatigue about 2 to 3 weeks into your treatment, and it may last for weeks to months after your treatment ends. Other treatments you're getting can affect how long fatigue lasts.
- **Nausea or vomiting:** Nausea is when you feel like you might vomit (throw up). You may have nausea, vomiting, or both during your treatment. We may prescribe an anti-nausea medication to help prevent this. You'll take the medication 30 minutes before each of your treatments.

Simulation date:

---

Treatment start date:

---

Medication instructions:

---

---

---

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

---

About Palliative Whole Brain Radiation Therapy - Last updated on June 1, 2026

All rights owned and reserved by Memorial Sloan Kettering Cancer Center