

PATIENT & CAREGIVER EDUCATION

Pelvic Floor Muscle (Kegel) Exercises for Females

This information explains what female pelvic floor muscle exercises (Kegels) are and how to do them.

About Kegel Exercises

Doing Kegel exercises can help strengthen your pelvic floor muscles. These muscles support your uterus, bladder, and bowel (see Figure 1). These are also called your pelvic organs.




Figure 1. Pelvic floor muscles and pelvic organs

How Kegel Exercises Can Help You

Kegel exercises can help you:

- Manage or prevent incontinence. Incontinence is when your urine (pee) or stool (poop) leak and you cannot control it.
- Support your pelvic organs. This helps decrease incontinence and pain.
- Improve your pelvic floor muscle coordination. This
 helps your muscles become strong and flexible. This is
 helpful if you have pain during sexual intercourse, pelvic
 exams, or both.

Kegel exercises can also improve your sexual health and

pleasure by:

- Improving blood circulation to your vagina and pelvic floor.
- Making it easier for you to reach orgasm.
- Increasing vaginal lubrication (wetness).

How to Find Your Pelvic Floor Muscles

It's important to find your pelvic floor muscles when doing Kegel exercises to make sure you're working out the right muscles.

To feel your pelvic floor muscles, try:

- Imagining you're urinating (peeing). Tighten the muscles you would use to stop the stream of urine. Those are your pelvic floor muscles.
 - Do not stop your urine stream when you urinate. This can keep your bladder from emptying fully. This can also increase your risk for a urinary tract infection (UTI).
- Tightening the muscles you would use to hold back a bowel movement or keep yourself from passing gas.
- Placing the tip of a clean finger into your vagina. Tighten your pelvic floor muscles around your finger. You should feel your vagina tighten and your pelvic floor move

upward.

How to Know If You're Using the Wrong Muscles

Do not tighten your abdomen (belly), leg, or buttock (butt) muscles when you contract your pelvic floor muscles.

- To find out if you're also contracting your leg or buttock muscles, place one hand underneath your buttocks or on the inside of your upper leg. Squeeze your pelvic floor muscles. If you feel your leg or buttocks move, you're using the wrong muscles.
- If you're sitting in a chair, your body should not lift from the chair when doing Kegel exercises. If you notice that your body lifts slightly, you're using your buttock muscles.

How to Do Kegel Exercises

Get Comfortable Before You Start

Before you start your Kegel exercises, get into a comfortable position so your body is relaxed. Most people prefer doing Kegel exercises while lying down on a bed or sitting in a chair. You can do them in any position you feel comfortable in.

Once you're familiar with the exercises, you should be able to do them in any position and in any place. This includes standing and waiting in a line, which can be a good way to

do them more often.

Follow These Steps to Do Kegels

- 1. Breathe in deeply through your nose and let your abdomen expand as it fills with air. Keep your pelvic floor muscles relaxed as you breathe in.
- 2. Breathe out slowly through your mouth as if you're blowing through a straw. Do this as you gently tighten your pelvic floor muscles.
- 3. Keep your pelvic floor muscles tightened for 3 to 6 seconds while you breathe out. This is called a contraction.
- 4. Breathe in again and release the contraction. This relaxes your muscles.
- 5. Relax your muscles completely for 6 to 10 seconds. It's very important that you relax your muscles fully between each contraction.

Do not hold your breath when doing Kegel exercises. If you feel yourself holding your breath, then count out loud when doing Kegel exercises.

How Often to Do Kegels

Repeat this exercise 10 times a session. Do 2 to 3 sessions of Kegel exercises every day to get the best results. Spread

the sessions out during the day.

When to Hold Your Kegels for Longer

You can start to hold your Kegels for longer if:

- Your pelvic floor muscles don't start to get tired after contracting them for 3 to 6 seconds.
- Your pelvic floor muscles don't start to get tired after doing 10 Kegel exercises in a row.

If this happens, try holding your contractions for 6 to 10 seconds. Then, relax your muscles completely for 10 seconds. Make sure you keep breathing while you hold the contractions.

Your goal should be to hold a strong contraction for 10 seconds 10 times in a row.

What to Do If You Have Pain from Kegel Exercises

Kegel exercises should not hurt. If you have pain during or after Kegel exercises, stop doing the exercises right away and call your healthcare provider. You may not be doing the exercise correctly or they may not be right for you.

Getting Sexual Health and Intimacy Support at MSK

If you need more support or information about sexual health or intimacy, talk with your healthcare provider. You

can also make an appointment with MSK's Female Sexual Medicine and Women's Health Program. For more information or to make an appointment, call 646-888-5076.

The Female Sexual Medicine and Women's Health Program provides services at the following locations:

- Rockefeller Outpatient Pavilion
 160 E. 53rd St.
 New York, NY 10022
- Evelyn H. Lauder Breast Center
 300 E. 66th St.
 New York, NY 10065

When to Call Your Healthcare Provider

Call your healthcare provider if you:

- Have concerns about your bowel, bladder, or sexual function.
- Are having trouble feeling your pelvic floor muscles.
- Have pain when you do Kegel exercises.
- Have trouble doing Kegel exercises.
- Have pelvic pain.
- Want a referral to a physical therapist who specializes in pelvic health.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Pelvic Floor Muscle (Kegel) Exercises for Females - Last updated on December 15, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center