



PATIENT & CAREGIVER EDUCATION

Pentobarbital

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Nembutal

What is this drug used for?

- It is used to treat sleep problems.
- It is used to calm a child before a procedure.
- It is used to treat seizures.
- It is used to put your child to sleep for surgery.
- It may be given to your child for other reasons. Talk with the doctor.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has ever had porphyria.
- If your child has liver disease.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for your child to take this drug with all of his/her drugs and

health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child avoid tasks or actions that call for alertness until the effects of this drug wear off and your child feels fully awake. These are things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized vehicles.
- Have your child's blood work checked if he/she is on this drug for a long time. Talk with your child's doctor.
- This drug may be habit-forming with long-term use.
- Give this drug for short periods of time. If signs show up again, talk with the doctor.
- If your child has been taking this drug on a regular basis and stops taking it all of a sudden, your child may have signs of withdrawal. Do not stop giving this drug all of a sudden without calling the doctor. Tell the doctor if your child has any bad effects.
- Alcohol may interact with this drug. Be sure your child does not drink alcohol.
- Talk with your child's doctor before giving your child other drugs and natural products that may slow your child's actions.
- Studies in young animals and children have shown that frequent or long-term use of anesthesia drugs or drugs used for sleep in children younger than 3 years of age may lead to long-term brain problems. This may also happen in unborn babies if the mother uses this drug during the third trimester of pregnancy. Talk with the doctor.

If your child is or may be sexually active:

- Birth control pills and other hormone-based birth control may not work as well to prevent pregnancy. Be sure your child uses some other kind of birth

control also, like a condom, when taking this drug.

If your child is pregnant or breast-feeding a baby:

- This drug may cause harm to the unborn baby if your child takes it during pregnancy. If your child is pregnant or gets pregnant while taking this drug, call the doctor right away.
- Long-term use of this drug during pregnancy may cause dependence in the unborn baby or newborn. Talk with the doctor.
- Tell the doctor if your child is breast-feeding a baby. You will need to talk about any risks to the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

For all uses of this drug:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Trouble breathing, slow breathing, or shallow breathing.
- Very bad dizziness or passing out.
- Feeling nervous and excitable.
- Feeling confused.
- Low mood (depression).

Injection (if given in the muscle):

- Pain and irritation where this drug goes into the body.

Injection (if given in the vein):

- This drug may cause tissue damage if the drug leaks from the vein. Tell your child's nurse if your child has any redness, burning, pain, swelling, blisters, skin sores, or leaking of fluid where the drug is going into your child's body.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Feeling sleepy.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into a muscle or vein.

What do I do if my child misses a dose?

- Call your child's doctor to find out what to do.

How do I store and/or throw out this drug?

- If you need to store this drug at home, talk with your child's doctor, nurse, or pharmacist about how to store it.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and

pets.

- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of using this medicine.

Last Reviewed Date

2019-03-19

Copyright

© 2019 Wolters Kluwer Clinical Drug Information, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your healthcare team directly. If you're a patient at MSK and you need to reach a provider after 5:00 PM, during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Pentobarbital - Last updated on November 18, 2019

©2019 Memorial Sloan Kettering Cancer Center