
PATIENT & CAREGIVER EDUCATION

Phenytoin

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Dilantin; Dilantin Infatabs; Phenytek; Phenytoin Infatabs

Brand Names: Canada

APO-Phenytoin Sodium; Dilantin; Dilantin Infatabs; Dilantin-125; Dilantin-30; NOVO-Phenytoin [DSC]; TARO-Phenytoin; Tremytoine

Warning

Injection:

- This drug may cause an abnormal heartbeat or low blood pressure if given too fast. Sometimes, this has been deadly. The doctor will watch your child's heart closely when your child gets this drug. Tell your child's doctor right away if your child has a fast, slow, or abnormal heartbeat during or after a dose.

What is this drug used for?

- It is used to help control certain kinds of seizures.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child is taking delavirdine.
- If your child has had liver problems in the past while taking this drug.
- If your child has a slow heartbeat, heart block, or other heartbeat that is not normal.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child's health care providers that your child is

taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.

- Have your child wear disease medical alert ID (identification).
- Have your child's blood work checked often. Talk with your child's doctor.
- Have the level of phenytoin in your child's blood checked as you have been told by the doctor. If the level is too high, some side effects may happen. This includes a type of brain problem that may not go back to normal. Talk with the doctor.
- Do not stop giving this drug to your child all of a sudden without calling the doctor. Your child may have a greater risk of seizures. If your child needs to stop this drug, you will want to slowly stop it as told by the doctor.
- Have your child avoid tasks or actions that call for alertness until you see how this drug affects your child. These are things like riding a bike, playing sports, or using items such as scissors, lawnmowers, electric scooters, toy cars, or motorized vehicles.
- This drug interacts with many other drugs. The chance of this drug's side effects may be raised or how well this drug works may be lowered. The chance of the other drugs' side effects may also be raised. This may include very bad, life-threatening, or deadly side effects. Check with the doctor and pharmacist to make sure that it is safe for your child to take

this drug with all of their other drugs (prescription or OTC, natural products, vitamins).

- People with a certain gene (HLA-B*1502) have a higher chance of severe and sometimes deadly skin reactions with this drug. This gene is more common in Asian people. If you have questions, talk with the doctor.
- This drug may cause weak bones. This may happen more often if used for a long time. This may raise the chance of broken bones. Call your doctor right away if you have bone pain.
- This drug may affect certain lab tests. Tell all of your child's health care providers and lab workers that your child takes this drug.
- If your child has high blood sugar (diabetes), talk with the doctor. This drug can raise blood sugar.
- Alcohol may interact with this drug. Be sure your child does not drink alcohol.
- Talk with your child's doctor before your child uses marijuana, other forms of cannabis, or prescription or OTC drugs that may slow your child's actions.
- Take care of your child's teeth. See a dentist often.

If your child is or may be sexually active:

- Birth control pills and other hormone-based birth control may not work as well to prevent pregnancy. Be sure your child uses

some other kind of birth control also, like a condom, when taking this drug.

If your child is pregnant or breast-feeding a baby:

- This drug may cause harm to the unborn baby if your child takes it during pregnancy. If your child is pregnant or gets pregnant while taking this drug, call the doctor right away.
- A bleeding problem that can be life-threatening may happen in newborns if your child took this drug during pregnancy.
- Tell the doctor if your child is breast-feeding a baby. You will need to talk about any risks to the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

All products:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.

- Signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- Trouble walking.
- Feeling confused.
- Slurred speech.
- Swollen or sore gums.
- Very bad muscle pain.
- Trouble controlling body movements, twitching, change in balance, trouble swallowing or speaking.
- A burning, numbness, or tingling feeling that is not normal.
- Not able to control eye movements.
- Shakiness.
- If seizures are different or worse after starting this drug, talk with the doctor.
- Heart problems like slow heartbeat have happened with this drug. Call your child's doctor right away if your child has any signs of heart problems like dizziness or passing out, chest pain or pressure, tiredness, or abnormal heartbeat.
- A severe skin reaction (Stevens-Johnson syndrome/toxic epidermal necrolysis) may happen. It can cause severe health problems that may not go away, and sometimes death. Get

medical help right away if your child has signs like red, swollen, blistered, or peeling skin (with or without fever); red or irritated eyes; or sores in the mouth, throat, nose, or eyes.

- Like other drugs that may be used for seizures, this drug may rarely raise the risk of suicidal thoughts or actions. The risk may be higher in people who have had suicidal thoughts or actions in the past. Call the doctor right away about any new or worse signs like depression; feeling nervous, restless, or grouchy; panic attacks; or other changes in mood or behavior. Call the doctor right away if any suicidal thoughts or actions occur.
- A very bad and sometimes deadly effect has happened in people taking drugs for seizures like this drug. Call the doctor right away if your child has swollen glands; fever; rash; chest pain; not able to pass urine or change in how much urine is passed; or signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Lymph node problems like cancer have happened in people taking this drug. It is not known if this drug may be the cause. Talk with the doctor if your child has swollen glands.
- Rarely, certain blood problems have happened with this drug. This can lead to bleeding problems or infections. Sometimes, these have been deadly. Tell the doctor right away if your child has signs of infection like fever, chills, very bad sore throat,

ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or a wound that will not heal. Tell the doctor right away if your child has any unexplained bruising or bleeding, or feels very tired or weak.

Injection:

- Change in skin color to black or purple.
- This drug may cause tissue damage if the drug leaks from the vein. Tell your child's nurse if your child has any redness, burning, pain, swelling, blisters, skin sores, or leaking of fluid where the drug is going into your child's body.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Feeling dizzy or sleepy.
- Headache.
- Feeling nervous and excitable.
- Trouble sleeping.
- Upset stomach or throwing up.
- Constipation.
- Change in taste.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

All oral products:

- Give this drug at the same time of day.
- Keep giving this drug to your child as you have been told by your child's doctor or other health care provider, even if your child feels well.
- Do not give antacids at the same time as this drug. Ask the doctor if you have a question about how to give antacids with this drug.
- If this drug is given through a feeding tube, stop tube feeding for 2 hours before giving this drug. Restart tube feeding 2 hours after giving.

Extended-release capsules:

- Have your child swallow whole. Do not let your child chew, break, or crush.

- Do not give if product changes color.

Chewable tablet:

- Have your child chew or swallow tablet whole.

Suspension:

- Shake well before use.
- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.
- Do not use a household teaspoon or tablespoon to measure this drug. Doing so could lead to the dose being too high.

Injection:

- It is given into a vein for a period of time.

What do I do if my child misses a dose?

All oral products:

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

Injection:

- Call your child's doctor to find out what to do.

How do I store and/or throw out this drug?

All oral products:

- Store at room temperature protected from light. Store in a dry place. Do not store in a bathroom.

Suspension:

- Do not freeze.

Injection:

- If you need to store this drug at home, talk with your child's doctor, nurse, or pharmacist about how to store it.

All products:

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better

or if they become worse, call your child's doctor.

- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- This drug comes with an extra patient fact sheet called a Medication Guide. Read it with care. Read it again each time this drug is refilled. If you have any questions about this drug, please talk with the doctor, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions,

warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of using this medicine.

Last Reviewed Date

2020-10-28

Copyright

© 2020 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your healthcare team directly. If you're a patient at MSK and you need to reach a provider after 5:00 PM, during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Phenytoin - Last updated on January 12, 2021

All rights owned and reserved by Memorial Sloan Kettering Cancer Center