



---

PATIENT & CAREGIVER EDUCATION

# Preventing An Allergic Reaction to Contrast Dye

This information is for people who have had an allergic reaction to contrast dye in the past. It explains how to prevent an allergic reaction to the contrast dye you'll get during your scan.

## About contrast dye

Contrast is a special dye used in imaging scans. It makes the images from the scans clearer and helps your doctors see your organs better.

There are different kinds of contrast. Computed tomography (CT) scans and magnetic resonance imaging (MRI) use different kinds of contrast. Having an allergic reaction to one type of contrast does not mean you'll need medicine before getting another type of contrast.

You've had an allergic reaction to contrast and you're having a scan that needs the same type of contrast. Because you've had an allergic reaction, you'll need to take a steroid medicine and an antihistamine (AN-tee-

HIS-tuh-meem) before your scan. This will help prevent another allergic reaction.

You're scheduled to have a \_\_\_\_\_ scan on \_\_\_\_\_ (date) at \_\_\_\_\_ AM | PM.

## **Preventing an allergic reaction to contrast with medicine**

There are different medicines you can take to prevent an allergic reaction to contrast. Your doctor will give you a prescription for the medicine you need.

Here are some medicines used to help prevent an allergic reaction to contrast. Ask your nurse if you have any questions about when to take your medicine.

### **Prednisone**

If your doctor gave you prednisone (PRED-nih-sone), take 3 doses before your scan:

- Take the first dose 13 hours before your scan.
- Take the second dose 7 hours before your scan.
- Take the third dose 1 hour before your scan.

Write down the time you need to take your medicine below.

Take 1 (50 milligram) tablet of prednisone at:

\_\_\_\_\_ AM | PM (13 hours before your scan) on \_\_\_\_\_  
(date)

\_\_\_\_\_ AM | PM (7 hours before your scan) on \_\_\_\_\_  
(date)

\_\_\_\_\_ AM | PM (1 hour before your scan) on \_\_\_\_\_  
(date)

Take 50 milligrams (mg) of diphenhydramine (Benadryl) with your last dose of prednisone 1 hour before your scan. If you're allergic to Benadryl, or do not want to be drowsy, take 10 mg of cetirizine (Zyrtec) instead.

## **Methylprednisolone**

If your doctor gave you methylprednisolone (MEH-thul-pred-NIH-suh-lone), take 2 doses before your scan:

- Take the first dose 12 hours before your scan.
- Take the second dose 2 hours before your scan.

Write down the time you need to take your medicine below.

Take 1 (32 milligram) tablet of methylprednisolone at:

\_\_\_\_\_ AM | PM (12 hours before your scan) on \_\_\_\_\_  
(date)

\_\_\_\_\_ AM | PM (2 hours before your scan) on \_\_\_\_\_

(date)

You may also need to take 50 mg of diphenhydramine 1 hour before your scan. Your doctor will tell you if you need to do this. If you're allergic to Benadryl, or do not want to be drowsy, take 10 mg of cetirizine (Zyrtec) instead.

## **Other medicine**

You may need to take more medicine, or a different medicine instead. Follow your doctor's instructions.

If you have any questions about your premedication plan, call the doctor that ordered your premedication.

Use the space below to write down your doctor's instructions and any other notes.

Take: \_\_\_\_\_ at \_\_\_\_\_ AM | PM, 1 hour before your scan.

## **Notes**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

If you have any questions, contact a member of your care team directly. If you're a patient at MSK and you need to reach a provider after 5 p.m., during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](http://www.mskcc.org/pe) to search our virtual library.

---

Preventing An Allergic Reaction to Contrast Dye - Last updated on January 4, 2024

All rights owned and reserved by Memorial Sloan Kettering Cancer Center