



PATIENT & CAREGIVER EDUCATION

Rabies Immune Globulin (Human)

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

HyperRAB; HyperRAB S/D [DSC]; Imogam Rabies-HT; Kedrab

Brand Names: Canada

HyperRAB; HyperRAB S/D [DSC]; Imogam Rabies Pasteurized [DSC]; Kamrab

What is this drug used for?

- It is used to prevent rabies.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your

doctor about the allergy and what signs you had.

- If you have started rabies vaccine treatment and already had a dose of this drug.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Talk with your doctor before getting any vaccines after getting this drug. Vaccine use after getting this drug may make the vaccine not work as well.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take

this drug.

- This drug is made from human plasma (part of the blood) and may have viruses that may cause disease. This drug is screened, tested, and treated to lower the chance that it carries an infection. Talk with the doctor.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of certain infections (parvovirus B19, hepatitis

A) like fever or chills, feeling very sleepy, runny nose, rash, joint pain, tiredness, poor appetite, upset stomach or throwing up, belly pain, or yellow skin or eyes.

- A burning, numbness, or tingling feeling that is not normal.
- Flushing.
- Fast breathing.
- Sweating a lot.
- Dark urine.
- There is a chance of blood clots with this drug. The chance is raised in older people and in people with thick blood, heart problems, or a history of blood clots. The chance may also be higher if you must be in a bed or chair for some time, if you take estrogen, or if you use certain catheters. However, blood clots can happen even if you do not have any of these health problems. Call your doctor right away if you have numbness or weakness on 1 side of your body; pain, redness, tenderness, warmth, or swelling in the arms or legs; change in color of an arm or leg; chest pain or pressure; shortness of breath; fast heartbeat; or coughing up blood. Talk with your doctor.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Pain where the shot was given.
- Irritation where the shot is given.
- Bruising.
- Headache.
- Stomach pain or diarrhea.
- Gas.
- Stuffy nose.
- Throat pain.
- Muscle or joint pain.
- Pain in arms or legs.
- Signs of a common cold.
- Feeling dizzy, tired, or weak.
- Throwing up.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into a muscle or into the skin.

What do I do if I miss a dose?

- This drug will be given in a health care setting.

How do I store and/or throw out this drug?

- This injection will be given to you in a healthcare setting. You will not store it at home.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Keep all drugs in a safe place. Keep all drugs out of

the reach of children and pets.

- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess

potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2024-02-12

Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Rabies Immune Globulin (Human - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center