



PATIENT & CAREGIVER EDUCATION

Somatropin

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Genotropin; Genotropin MiniQuick; Humatrope; Norditropin FlexPro; Nutropin AQ NuSpin 10; Nutropin AQ NuSpin 20; Nutropin AQ NuSpin 5; Nutropin AQ Pen [DSC]; Omnitrope; Saizen; Saizen Click.Easy [DSC]; Saizenprep; Serostim; Zomacton; Zorbtive

Brand Names: Canada

Genotropin GoQuick; Genotropin MiniQuick; Humatrope; Norditropin NordiFlex Pen; Nutropin AQ NuSpin 10; Nutropin AQ NuSpin 20; Nutropin AQ NuSpin 5; Nutropin AQ Pen [DSC]; Omnitrope; Saizen; Serostim

What is this drug used for?

- It is used to help with growth and to treat growth hormone deficiency.
- It may be given to your child for other reasons. Talk with the doctor.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child has an allergy to any part of this drug.
- If your child is allergic to any drugs like this one or any other drugs, foods, or other substances. Tell the doctor about the allergy and what signs your child had, like rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other signs.
- If your child has any of these health problems: Breathing problems like sleep apnea; cancer or other tumors like a brain tumor; diabetic eye disease; or illness shortly after open heart surgery, stomach surgery, or accidental injury.
- If your child has Prader-Willi syndrome and is very overweight, has trouble breathing, or has sleep apnea.
- If your child's bones are no longer growing (closed epiphyses).

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for your child to take this drug with all of his/her drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my

child takes this drug?

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- High blood sugar has happened with this drug. This includes diabetes that is new or worse. Talk with the doctor.
- Have your child's blood sugar checked as you have been told by your child's doctor.
- This drug may affect certain lab tests. Tell all of your child's health care providers and lab workers that your child takes this drug.
- Have your child's blood work checked often. Talk with your child's doctor.
- Have your child's eye pressure and eyesight checked as you have been told by the doctor. Talk with the doctor.
- If your child has an abnormal curve in the spine (scoliosis), talk with your doctor. This drug could make it worse in children who are still growing.
- If your child has Turner syndrome, talk with the doctor. The chance of ear infections, high blood pressure, and very bad blood vessel problems like stroke and bleeding in the brain may be raised.
- If your child has cancer or a tumor or has ever had cancer or a tumor, talk with your child's doctor. The chance of cancer or tumor growth is raised with this drug. The chance of new

tumors may also be raised in some patients.

- Some products have benzyl alcohol. Do not give a product that has benzyl alcohol in it to a newborn or infant. Talk with the doctor to see if this product has benzyl alcohol in it.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain, or very bad upset stomach

or throwing up.

- Signs of a weak adrenal gland like a very bad upset stomach or throwing up, very bad dizziness or passing out, muscle weakness, feeling very tired, mood changes, not hungry, or weight loss.
- Signs of low thyroid levels like constipation; not able to handle cold; memory problems; mood changes; or a burning, numbness, or tingling feeling that is not normal.
- Signs of a urinary tract infection (UTI) like blood in the urine, burning or pain when passing urine, feeling the need to pass urine often or right away, fever, lower stomach pain, or pelvic pain.
- Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight.
- Shortness of breath, a big weight gain, or swelling in the arms or legs.
- Chest pain or pressure or a fast heartbeat.
- Sweating a lot.
- Mood changes.
- Low mood (depression).
- Change in how you act.
- Change in skin color.
- Burning, numbness, pain, or tingling in the hands, arms,

legs, or feet.

- Bone pain.
- Change in color or size of a mole.
- Redness or swelling where the shot is given.
- Skin breakdown where this drug is used.
- Children who use this drug can rarely have a bone problem in the hip (slipped growth plate). Call the doctor right away if your child has hip or knee pain or a limp.
- Raised pressure in the head has rarely happened with this drug. The risk may be greater in patients with Turner syndrome or Prader-Willi syndrome. Signs most often happen within the first 8 weeks of starting this drug. Call your child's doctor right away if your child has a change in eyesight, a severe headache, upset stomach, or throwing up.
- Very bad and sometimes deadly lung or breathing problems have happened in some children with Prader-Willi syndrome. The chance may be higher in children who have sleep apnea, an infection in the lungs or airway, a block in the airway, and in children who are very overweight. Call the doctor right away if your child has a fever of 100.4°F (38°C) or higher, cough, sore throat, shortness of breath, chest pain or discomfort, new or worse snoring, or breathing that is not normal while asleep.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have

no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Headache.
- Feeling tired or weak.
- Muscle or joint pain.
- Muscle stiffness.
- Back pain.
- Trouble sleeping.
- Upset stomach or throwing up.
- Gas.
- Stomach pain or diarrhea.
- Irritation where the shot is given.
- Signs of a common cold.
- Hair loss.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- It is given as a shot into the fatty part of the skin.
- If you will be giving your child the shot, your child's doctor or nurse will teach you how to give the shot.
- Move the site where you give the shot with each shot.
- Do not give into skin that is irritated, bruised, red, infected, or scarred.
- Wash your hands before and after use.
- Do not shake the solution.
- Do not use if the solution is cloudy, leaking, or has particles.
- Do not use if solution changes color.
- Throw away needles in a needle/sharp disposal box. Do not reuse needles or other items. When the box is full, follow all local rules for getting rid of it. Talk with a doctor or pharmacist if you have any questions.

What do I do if my child misses a dose?

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child's next dose, skip the missed dose and go back to your child's normal time.
- Do not give 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Some brands of this drug need to be stored in a refrigerator. Some brands of this drug need to be stored at room temperature. If you have questions about how to store

this drug, talk with your pharmacist.

- Protect from light.
- Be sure you know how long you can store this drug before you need to throw it away.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Keep a list of all your child's drugs (prescription, natural products, vitamins, OTC) with you. Give this list to your child's doctor.
- Talk with your child's doctor before giving your child any new drug, including prescription or OTC, natural products, or vitamins.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your

child's doctor, nurse, pharmacist, or other health care provider.

- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This information should not be used to decide whether or not to take this medicine or any other medicine. Only the healthcare provider has the knowledge and training to decide which medicines are right for a specific patient. This information does not endorse any medicine as safe, effective, or approved for treating any patient or health condition. This is only a brief summary of general information about this medicine. It does NOT include all information about the possible uses, directions, warnings, precautions, interactions, adverse effects, or risks that may apply to this medicine. This information is not specific medical advice and does not replace information you receive from the healthcare provider. You must talk with the healthcare provider for complete information about the risks and benefits of using this medicine.

Last Reviewed Date

2019-06-12

Copyright

© 2019 Wolters Kluwer Clinical Drug Information, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your healthcare team directly. If you're a patient at MSK and you need to reach a provider after 5:00 PM, during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Somatropin - Last updated on October 21, 2019

©2019 Memorial Sloan Kettering Cancer Center