Memorial Sloan Kettering Cancer Center

PATIENT & CAREGIVER EDUCATION

Tenapanor

This information from Lexicomp® explains what you need to know about this medication, including what it’s used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Ibsrela; Xphozah

Brand Names: Canada

Ibsrela

Warning
When used to treat irritable bowel syndrome:

- In studies, some young rodents died from dehydration after taking this drug. Avoid use of this drug in children younger than 18 years of age. If you have questions, talk with the doctor.

What is this drug used for?

- It is used to treat irritable bowel syndrome with constipation (IBS-C).
- It is used to lower high phosphate levels.

What do I need to tell my doctor BEFORE I take this drug?

For all uses of this drug:

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have a bowel block.

When used to lower phosphate levels:

- If the patient is a child. This drug is not approved for use in children. Do not give this drug to a child younger than 6 years of age.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose

of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

For all uses of this drug:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.

When used to treat irritable bowel syndrome:

- Swelling of belly.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

For all uses of this drug:

- Diarrhea is common with this drug and can be severe. Call your doctor to find out what to do if you have diarrhea or stomach cramps. You will need to take care not to become dehydrated. Do not try to treat diarrhea without first checking with your doctor.

When used to treat irritable bowel syndrome:

- Stomach pain.
- Gas.
- Bloating.
- Dizziness.
- Nose or throat irritation.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor.

Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at https://www.fda.gov/medwatch.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

For all uses of this drug:

- Take the first dose of the day right before breakfast or the first meal of the day. Take the second dose of the day right before dinner. Take this way unless your doctor has told you something else.

When used to treat irritable bowel syndrome:

- Swallow whole. Do not chew, break, or crush.

When used to lower phosphate levels:

- Do not take this drug right before a dialysis session. Take it right before the next meal after dialysis. If you have questions about when to take this drug, talk with your doctor.

- If you are also taking sodium polystyrene sulfonate (SPS), take it at least 3 hours before or 3 hours after this drug.

What do I do if I miss a dose?

- Skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store tablets in the original container at room temperature. Keep the cap tightly closed. Do not take out the antimoisture cube or packet.
- Store in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.

- Do not share your drugs with others and do not take anyone else’s drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice

or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2023-11-07

Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your care team directly. If you're a patient at MSK and you need to reach a provider after 5 p.m., during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Tenapanor - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center