

About Your Hip Replacement Surgery

This guide will help you get ready for your hip replacement surgery at MSK. It will also help you know what to expect as you recover.

Use this guide as a source of information in the days leading up to your surgery. Bring it with you on the day of your surgery. You and your care team will use it as you learn more about your recovery.

Your Care Team

Doctor: _____

Nurse: _____

Phone number: _____

Fax number: _____

Your Caregiver

It's important to choose someone to help you care for yourself before and after your surgery. This person is called your caregiver.

Your caregiver will learn about your surgery with you. They'll also help you care for yourself while you're healing after surgery. Write their name below.

Caregiver: _____



Visit www.mskcc.org/pe/hip_replacement to view this guide online.

Table of Contents

About Your Hip Replacement Surgery.....	3
Your Hip Anatomy.....	3
Your Hip Replacement Surgery.....	4
Getting Ready for Your Hip Replacement Surgery.....	5
Getting Ready for Your Surgery	6
Within 30 Days of Your Hip Replacement Surgery	8
7 Days Before Your Hip Replacement Surgery.....	11
2 Days Before Your Hip Replacement Surgery.....	12
1 Day Before Your Hip Replacement Surgery.....	12
The Morning of Your Hip Replacement Surgery	15
Recovering After Your Hip Replacement Surgery	21
In the Post-Anesthesia Care Unit (PACU).....	22
In Your Hospital Room	23
When To Call Your Healthcare Provider.....	40
Support Services.....	41
MSK Support Services	42
External Support Services.....	46
Educational Resources	47
<i>How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil</i>	
<i>Herbal Remedies and Cancer Treatment</i>	
<i>How To Use Your Incentive Spirometer</i>	

About Your Hip Replacement Surgery

Your Hip Anatomy

Your hip is a ball and socket joint. This means it's made up of 2 parts that allow for movement and rotation. Your hip is made up of the femoral head, which is the top end of your femur (thigh bone). There also is a socket in your pelvis called the acetabulum (see Figure 1). The femoral head is a round surface that looks like a ball. It fits into the acetabulum and allows smooth motion of the hip.




Figure 1. Your hip anatomy

The parts of your hip joint that are damaged will be replaced with artificial parts, called prostheses. These are usually made of metal or plastic. The new joint will help reduce pain and increase your ability to move around.

Types of Hip Replacements

There are 2 types of hip replacements, a partial hip replacement and a total hip replacement.

In a partial hip replacement, the femoral head is removed and replaced with a metal stem inserted into your femur. A ball is placed on the top part of the stem. A device called a “bipolar cup” snaps on the ball of your new hip and rotates in your own hip socket.

In a total hip replacement, your femoral head is replaced, and your acetabulum is replaced with a metal socket. The metal socket is attached to your bone by cement, screws, or your bone itself.

Your Hip Replacement Surgery

Your surgeon will make an incision (surgical cut). It will run about 5 inches (12.7 centimeters) above your hip to about 6 inches (15.24 centimeters) below your hip. They'll remove the diseased portion of your bone and replace it with your prosthesis. It will be fitted and fixed in place.

Notes _____

Getting Ready for Your Hip Replacement Surgery

This section will help you get ready for your surgery. Read it when your surgery is scheduled. Refer to it as your surgery gets closer. It has important information about what to do to get ready.

As you read this section, write down questions to ask your healthcare provider. You can use the space below.

Notes _____

Getting Ready for Your Surgery

You and your care team will work together to get ready for your surgery. Help us keep you safe by telling us if any of the following statements apply to you, even if you're not sure.

☐ I take a blood thinner, such as:

- Aspirin
- Heparin
- Warfarin (Jantoven® or Coumadin®)
- Clopidogrel (Plavix®)
- Enoxaparin (Lovenox®)
- Dabigatran (Pradaxa®)
- Apixaban (Eliquis®)
- Rivaroxaban (Xarelto®)

There are others, so be sure your healthcare provider knows all the medications you're taking.

☐ I take prescription medications (medications my healthcare provider prescribes), including patches and creams.

☐ I take over-the-counter medications (medications I buy without a prescription), including patches and creams.

☐ I take dietary supplements, such as herbs, vitamins, minerals, or natural or home remedies.

☐ I have a pacemaker, automatic implantable cardioverter-defibrillator (AICD), or other heart device.

☐ I have sleep apnea.

☐ I've had a problem with anesthesia (medication to make me sleep during surgery) in the past.

☐ I'm allergic to certain medication(s) or materials, including latex.

☐ I'm not willing to receive a blood transfusion.

☐ I drink alcohol.

☐ I smoke or use an electronic smoking device, such as a vape pen, e-cigarette, or Juul®.

☐ I use recreational drugs.

About Drinking Alcohol

The amount of alcohol you drink can affect you during and after your surgery. It's important to talk with your healthcare providers about how much alcohol you drink. This will help us plan your care.

- If you stop drinking alcohol suddenly, it can cause seizures, delirium, and death. If we know you're at risk for these problems, we can prescribe medications to help keep them from happening.
- If you drink alcohol regularly, you may be at risk for other problems during and after your surgery. These include bleeding, infections, heart problems, and a longer hospital stay.

Here are things you can do before your surgery to keep from having problems:

- Be honest with your healthcare providers about how much alcohol you drink.
- Try to stop drinking alcohol once your surgery is planned. Tell your healthcare provider right away if you:
 - Get a headache.
 - Feel nauseous (like you're going to throw up).
 - Feel more anxious (nervous or worried) than usual.
 - Cannot sleep.

These are early signs of alcohol withdrawal and can be treated.

- Tell your healthcare provider if you cannot stop drinking.
- Ask your healthcare provider questions about drinking and surgery. As always, all your medical information will be kept private.

About Smoking

If you smoke, you can have breathing problems when you have surgery. Stopping for even a few days before surgery can help.

Your healthcare provider will refer you to our Tobacco Treatment Program if you smoke. You can also reach the program by calling 212-610-0507.

About Sleep Apnea

Sleep apnea is a common breathing problem. It causes you to stop breathing for short lengths of time while you're asleep. The most common type is obstructive sleep apnea (OSA). With OSA, your airway becomes fully blocked during sleep.

OSA can cause serious problems during and after surgery. Please tell us if you have or think you might have sleep apnea. If you use a breathing device (such as a CPAP machine), bring it on the day of your surgery.

Using MyMSK

MyMSK (my.mskcc.org) is your MSK patient portal. You can use it to send and read messages from your care team, view your test results, see your appointment dates and times, and more. You can also invite your caregiver to make their own account so they can see information about your care.

If you do not have a MyMSK account, you can sign up at my.mskcc.org. You can get an enrollment ID by calling 646-227-2593 or your doctor's office.

For help, watch *How to Enroll in MyMSK: Memorial Sloan Kettering's Patient Portal* at www.mskcc.org/pe/enroll_mymsk. You can also contact the MyMSK Help Desk by emailing mymsk@mskcc.org or calling 800-248-0593.

Within 30 Days of Your Hip Replacement Surgery

Presurgical Testing (PST)

You'll have a PST appointment before your surgery. The date, time, and location will be printed on the appointment reminder from your surgeon's office. You can eat and take your usual medications the day of your appointment.

It's helpful to bring these things to your appointment:

- A list of all the medications you're taking, including prescription and over-the-counter medications, patches, and creams.
- Results of any tests done outside of MSK, such as a cardiac stress test, echocardiogram, or carotid doppler study.
- The names and telephone numbers of your healthcare providers.

During your PST appointment, you'll meet with a nurse practitioner (NP). They work closely with anesthesiology staff (specialized healthcare providers who will give you anesthesia during your surgery). Your NP will review your medical and surgical history with you. You may have tests to plan your care. Examples are:

- An electrocardiogram (EKG) to check your heart rhythm.
- A chest X-ray.
- Blood tests.

Your NP may recommend you see other healthcare providers. They'll also talk with you about which medications to take the morning of your surgery.

Identify Your Caregiver

Your caregiver plays an important role in your care. Before your surgery, you and your caregiver will learn about your surgery from your healthcare providers. After your surgery, your caregiver will take you home when you're discharged. They'll also help you care for yourself at home.



For Caregivers

Caring for a person going through cancer treatment comes with many responsibilities. We offer resources and support to help you manage them.

For information, visit www.mskcc.org/caregivers or read *A Guide for Caregivers*. You can ask your healthcare provider for a copy or find it at www.mskcc.org/pe/guide_caregivers

Complete a Health Care Proxy Form

If you have not already filled out a Health Care Proxy form, we recommend you do now. If you already filled one out or have any other advance directives, bring them to your next appointment.

A health care proxy is a legal document. It says who will speak for you if you cannot communicate for yourself. This person is called your health care agent.

- For information about health care proxies and other advance directives, read *Advance Care Planning*. You can find it at www.mskcc.org/advance_care_planning or ask your healthcare provider for a copy.
- For information about being a health care agent, read *How to Be a Health Care Agent*. You can ask your healthcare provider for a copy or find it at www.mskcc.org/pe/health_care_agent
- If you have more questions about filling out a Health Care Proxy form, talk with your healthcare provider.

Do Breathing and Coughing Exercises

Practice taking deep breaths and coughing before your surgery. Your healthcare provider will give you an incentive spirometer to help expand your lungs.

For more information, read *How to Use Your Incentive Spirometer*. You can find it in the “Educational Resources” section of this guide.

Exercise

Exercising will help your body get into its best condition for your surgery and make your recovery faster and easier.

Try to do aerobic exercise every day. Aerobic exercise is any exercise that makes your heart beat faster, such as walking, swimming, or biking. If it's cold outside, use stairs in your home or go to a mall or shopping center.

Follow a Healthy Diet

Follow a well-balanced, healthy diet before your surgery. If you need help with your diet, talk with your healthcare provider about meeting with a clinical dietitian nutritionist.

Buy a 4% Chlorhexidine Gluconate (CHG) Solution Antiseptic Skin Cleanser (Such as Hibiclens®)

4% CHG solution is a skin cleanser that kills germs for 24 hours after you use it. Showering with it before your surgery will help lower your risk of infection after surgery. You can buy a 4% CHG solution antiseptic skin cleanser at your local pharmacy without a prescription.

7 Days Before Your Hip Replacement Surgery

Follow your healthcare provider's instructions for taking aspirin

Aspirin can cause bleeding. If you take aspirin or a medicine that has aspirin, you may need to change your dose or stop taking it 7 days before your surgery. Follow your healthcare provider's instructions. **Do not stop taking aspirin unless they tell you to.**

To learn more, read *How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil*. You can find it in the “Educational resources” section of this guide.

Stop taking vitamin E, multivitamins, herbal remedies, and other dietary supplements

Vitamin E, multivitamins, herbal remedies, and other dietary supplements can cause bleeding. Stop taking them 7 days before your surgery. If your healthcare provider gives you other instructions, follow those instead.

To learn more, read *Herbal Remedies and Cancer Treatment*. You can find it in the “Educational resources” section of this guide.

2 Days Before Your Hip Replacement Surgery

Stop taking nonsteroidal anti-inflammatory drugs (NSAIDs)

NSAIDs, such as ibuprofen (Advil® and Motrin®) and naproxen (Aleve®), can cause bleeding. Stop taking them 2 days before your surgery. If your healthcare provider gives you other instructions, follow those instead.

To learn more, read *How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil*. You can find it in the “Educational resources” section of this guide.

1 Day Before Your Hip Replacement Surgery

Note the Time of Your Surgery

A staff member from the Admitting Office will call you after 2 p.m. the day before your surgery. If your surgery is scheduled for a Monday, they'll call you the Friday before. If you do not get a call by 7 p.m., call 212-639-5014.

The staff member will tell you what time to arrive at the hospital for your surgery. They'll also remind you where to go.

This will be one of the following locations:

- Presurgical Center (PSC) on the 2nd floor
1275 York Avenue
(between East 67th and East 68th Streets)
New York, NY 10065
M Elevator to 2nd floor
- Presurgical Center (PSC) on the 6th floor
1275 York Avenue
(between East 67th and East 68th Streets)
New York, NY 10065
B elevator to 6th floor

Shower With a 4% CHG Solution Antiseptic Skin Cleanser (Such as Hibiclens)

The night before your surgery, shower with a 4% CHG solution antiseptic skin cleanser.

1. Wash your hair with your usual shampoo and conditioner. Rinse your head well.
2. Wash your face and genital (groin) area with your usual soap. Rinse your body well with warm water.
3. Open the 4% CHG solution bottle. Pour some into your hand or a clean washcloth.
4. Move away from the shower stream. Rub the 4% CHG solution gently over your body from your neck to your feet. Do not put it on your face or genital area.
5. Move back into the shower stream to rinse off the 4% CHG solution. Use warm water.
6. Dry yourself off with a clean towel.

Do not put on any lotion, cream, deodorant, makeup, powder, perfume, or cologne after your shower.

Notes _____

Instructions for eating and drinking: 8 hours before your arrival time



- **Stop eating 8 hours before your arrival time, if you have not already.**
 - Your healthcare provider may tell you to stop eating earlier. If they do, follow their instructions.
- 8 hours before your arrival time, do not eat or drink anything except these clear liquids:
 - Water.
 - Soda.
 - Clear juices, such as lemonade, apple, and cranberry juices. Do not drink orange juice or juices with pulp.
 - Black coffee or tea (without any type of milk or creamer).
 - Sports drinks, such as Gatorade®.
 - ClearFast CF(Preop)® or Ensure® Pre-Surgery clear carbohydrate drink.
 - Gelatin, such as Jell-O®.

You can keep having these until 2 hours before your arrival time.

The Morning of Your Hip Replacement Surgery

Remember, starting 8 hours before your arrival time, do not eat or drink anything except the things listed earlier.

Instructions for drinking: 2 hours before your arrival time



Stop drinking 2 hours before your arrival time. This includes water.

Take your medicines as instructed

A member of your care team will tell you which medicines to take the morning of your surgery. Take only those medicines with a sip of water. Depending on what you usually take, this may be all, some, or none of your usual morning medicines.

Shower With a 4% CHG Solution Antiseptic Skin Cleanser (Such as Hibiclens)

Shower with a 4% CHG solution antiseptic skin cleanser before you leave for the hospital. Use it the same way you did the night before.

Do not put on any lotion, cream, deodorant, makeup, powder, perfume, or cologne after your shower.

Things to Remember

- Remove nail polish and nail wraps.
- Wear something comfortable and loose-fitting.
- If you wear contact lenses, wear your glasses instead. Wearing contact lenses during surgery can damage your eyes.

- Before you're taken into the operating room, you'll need to remove your hearing aids, dentures, prosthetic device(s), wig, and religious articles.
- Do not wear any metal objects. Remove all jewelry, including body piercings. The tools used during your surgery can cause burns if they touch metal.
- Leave valuable items at home.
- If you're menstruating (have your monthly period), use a sanitary pad, not a tampon. You'll get disposable underwear, as well as a pad if needed.

What To Bring

- Sneakers that lace up or slip-on shoes with a sole. You may have some swelling in your feet. Lace-up sneakers or sturdy slip-on shoes will be easier to put on after surgery. Avoid open back shoes, such as flip flops or mules.
- Your breathing device for sleep apnea (such as your CPAP machine), if you have one.
- Your incentive spirometer, if you have one.
- Your Health Care Proxy form and other advance directives, if you filled them out.
- Your cell phone and charger.
- Only the money you may want for small purchases, such as a newspaper.
- A case for your personal items, if you have any. Examples of personal items include eyeglasses, hearing aids, dentures, prosthetic devices, wigs, and religious articles.
- This guide. You'll use it when you learn how to care for yourself after surgery.

Where To Park

MSK's parking garage is on East 66th Street between York and 1st avenues. If you have questions about prices, call 212-639-2338.

To reach the garage, turn onto East 66th Street from York Avenue. The garage is about a quarter of a block in from York Avenue. It's on the right (north) side of the street. There's a tunnel you can walk through that connects the garage to the hospital.

There are other parking garages located on:

- East 69th Street between 1st and 2nd avenues.
- East 67th Street between York and 1st avenues.
- East 65th Street between 1st and 2nd avenues.

Once You're in the Hospital

When you get to the hospital, take the B elevator to the 6th floor. Check in at the desk in the PSC waiting room.

Many staff members will ask you to say and spell your name and birth date. This is for your safety. People with the same or a similar name may be having surgery on the same day.

When it's time to change for surgery, you'll get a hospital gown, robe, and nonskid socks to wear.



For Caregivers, Family, and Friends

Read *Information for Family and Friends for the Day of Surgery* to help you know what to expect on the day of your loved one's surgery. You can or ask for a printed copy or find it at www.mskcc.org/pe/info_family_friends

Meet With a Nurse

You'll meet with a nurse before surgery. Tell them the dose of any medications you took after midnight and the time you took them. Make sure to include prescription and over-the-counter medications, patches, and creams.

Your nurse may place an intravenous (IV) line in one of your veins, usually in your arm or hand. If your nurse does not place the IV, your anesthesiologist will do it in the operating room.

Meet With an Anesthesiologist

You'll also meet with an anesthesiologist before surgery. They will:

- Review your medical history with you.
- Ask if you've had any problems with anesthesia in the past, including nausea or pain.
- Talk with you about your comfort and safety during your surgery.
- Talk with you about the kind of anesthesia you'll get.
- Answer your questions about your anesthesia.

Your doctor or anesthesiologist may also talk with you about placing an epidural catheter (thin, flexible tube) in your spine (back). An epidural catheter is another way to give you pain medication after your surgery.

You may also receive your pain medication through:

- A nerve catheter placed near the nerves that cause pain from your incision.
- Your IV.

Get Ready for Surgery

When it's time for your surgery, you'll need to remove your eyeglasses, hearing aids, dentures, prosthetic devices, wig, and religious articles.

You'll either walk into the operating room or a staff member will bring you there on a stretcher. A member of the operating room team will help you onto the operating bed. They'll place compression boots on your lower legs. These gently inflate and deflate to help blood flow in your legs.

Once you're comfortable, your anesthesiologist will give you anesthesia through your IV line and you'll fall asleep. You'll also get fluids through your IV line during and after your surgery.

During Your Surgery

After you're fully asleep, your care team will place a breathing tube through your mouth into your airway. It will help you breathe. They'll also place a urinary (Foley) catheter in your bladder. It will drain your urine (pee) during your surgery. Once they finish your surgery, your surgeon will close your incisions with staples or stitches. They may also place Steri-Strips™ (thin pieces of surgical tape) or Dermabond® (surgical glue) over your incisions. They'll cover your incisions with a bandage.

Your breathing tube is usually taken out while you're still in the operating room.

Notes _____

Recovering After Your Hip Replacement Surgery

This section will help you know what to expect after your surgery. You'll learn how to safely recover from your surgery both in the hospital and at home.

As you read this section, write down questions to ask your healthcare provider. You can use the space below.

Notes _____

In the Post-Anesthesia Care Unit (PACU)

When you wake up after your surgery, you'll be in the PACU. A nurse will be keeping track of your temperature, pulse, blood pressure, and oxygen levels. You may get oxygen through a tube resting below your nose or a mask over your nose and mouth. You'll also have compression boots on your lower legs.

Pain Medication

You'll get epidural or IV pain medication while you're in the PACU.

- If you're getting epidural pain medication, it will be put into your epidural space through your epidural catheter. Your epidural space is the space in your spine just outside your spinal cord.
- If you're getting IV pain medication, it will be put into your bloodstream through your IV line.

You'll be able to control your pain medication using a button called a patient-controlled analgesia (PCA) device. For more information, read *Patient-Controlled Analgesia (PCA)*. You can or ask your healthcare provider for a copy or find it at www.mskcc.org/pe/pca.

Tubes and Drains During your surgery, plastic drainage tubes will be placed in the area of the implant and will exit near your incision. These tubes will drain fluid from the surgical area to help reduce swelling. The drainage tubes may be stitched in place so they don't fall out. They're connected to a container that collects the drainage.

Moving to Your Hospital Room

You may stay in the PACU for a few hours or overnight. How long you stay depends on the type of surgery you had. After your stay in the PACU, a staff member will bring you to your hospital room.

In Your Hospital Room

The length of time you're in the hospital after your surgery depends on your recovery. Most people stay in the hospital for 4 days.

In your hospital room, you'll meet one of the nurses who will care for you during your stay. Soon after you get there, a nurse will help you out of bed and into your chair. Your healthcare providers will teach you how to care for yourself while you're healing from your surgery.

Managing Your Pain

You'll have some pain from your incisions after your surgery. At first, you'll get your pain medication through your epidural catheter or IV line. You'll be able to control your pain medication using a PCA device. Once you're able to eat, you'll get oral pain medication (medication you swallow).

Your healthcare providers will ask you about your pain often and give you medication as needed. If your pain is not relieved, tell one of your healthcare providers. It's important to control your pain so you can use your incentive spirometer and move around. Controlling your pain can help you recover faster.

You'll get a prescription for pain medication before you leave the hospital. Talk with your healthcare provider about possible side effects. Ask them when to start switching to over-the-counter pain medications.

The skin below your incision will feel numb after your surgery. This is because some of your nerves were cut. The numbness will go away over time.

Moving Around and Walking

Moving around and walking will help lower your risk for blood clots and pneumonia (lung infection). It will also help you start passing gas and having bowel movements (pooping) again. Your nurse, physical therapist, or occupational therapist will help you move around, if needed.

Read *Frequently Asked Questions About Walking After Your Surgery* to learn more about how walking after surgery can help you recover. You can find it at www.mskcc.org/pe/walking_after_surgery or ask your healthcare provider for a copy.

Read *Call! Don't Fall!* to learn what you can do to stay safe and keep from falling while you're in the hospital. You can ask a member of your care team for a copy or find it at www.mskcc.org/pe/call_dont_fall

Leg exercises can help to promote good circulation and prevent blood clots. You can find them in the “Exercises To Do While You’re in The Hospital” section of this resource. Do these exercises 10 times an hour while you’re awake.

After your hip replacement surgery, you must follow special hip precautions.

Depending on the surgery you had, these precautions may vary slightly. Your physical therapist (PT) and occupational therapist (OT) will go over these precautions with you. Most people will need to follow the precautions below:

- Do not bend at the waist or hip more than 90 degrees.
- Do not cross your legs while standing, sitting, or lying down.
- Do not rotate your operated leg (leg you had surgery on) inward.
- Do not lie, sit, or stand with “pigeon-toes.”

If you have any additional or different precautions, your doctor, PT, and OT will review these with you. During your stay in the hospital, your PT and OT will teach you how to move safely in and out of bed. They’ll also show you how to complete your daily self-care tasks, such as getting dressed and going to the bathroom, while following these precautions. For more information, read the “Hip Precautions” section of this resource.

Exercises To Do While You're in the Hospital

Ankle Pumps




Figure 1. Ankle pumps

1. Lie on your back with your head and shoulders supported on pillows. You can also do this exercise while sitting.
2. Point your toes up toward your nose (see Figure 1). You can do this with both feet at the same time.
3. Then, point them toward the floor.
4. Repeat 10 times.

Do this exercise every hour that you're awake.

Quadriceps Sets




Figure 2. Quadriceps sets

1. Lie on your back with your head and shoulders supported on pillows.
2. Straighten your legs as much as you can.
3. Push the backs of your knees down into the bed while tightening the muscles on the top of your thighs (see Figure 2).
4. Hold the position and count out loud to 5.
5. Relax.
6. Repeat 10 times.

Do this exercise every hour that you're awake.

Gluteal Sets



Figure 3. Gluteal sets

1. Lie on your back with your head and shoulders supported on pillows.
2. Straighten your legs as much as you can.
3. Squeeze your buttocks together tightly (see Figure 3).
4. Hold the position and count out loud to 5.
5. Relax your buttocks.
6. Repeat 10 times.

Do this exercise every hour that you are awake.

Hip Precautions

Safe moving during everyday activities

You need to follow special hip precautions. Here are some ways that you can adjust your activities to make sure that you follow your precautions.

Stand with your toes pointing straight ahead, about 6 to 10 inches (15 to 25 centimeters) apart. Do not stand with your toes pointing in or out (see Figure 1).




Figure 1. Standing positions after a hip replacement

When you need to pick something up, use your reacher (see Figure 2). Do not bend over.




Figure 2. Using reacher to pick up things

Move your upper and lower body as one unit. Make sure that your shoulders, hips, and knees are pointing in the same direction as your feet (see Figure 3). Do not twist your body.




Figure 3. Moving upper and lower body as one unit

Scoot to the edge and use the arms of your chair when rising from a chair (see Figure 4). Do not bend forward to stand up.




Figure 4. Standing up from a chair

Sit on a raised toilet seat or on a firm cushion in a chair (see Figure 5). Do not sit on low seats.




Figure 5. Sitting on firm chairs

Sit with your legs uncrossed (see Figure 6). Do not cross your legs while sitting or lying down.




Figure 6. Sitting with uncrossed legs

Put on your slippers when you're standing (see Figure 7). Do not lean down or over to the side to get your slippers from under the bed.




Figure 7. Putting on slippers while standing

Pull the blanket and sheets halfway up before getting into bed (see Figure 8). Do not lean forward to pull them up after getting into bed.




Figure 8. Pulling on blankets before getting into bed

Sleep with 1 pillow between your legs when lying on your back. Sleep with 2 pillows when lying on side. You can also lie on your stomach, but you must:

- Turn toward your unoperated side (side you didn't have surgery on).
- Have a pillow between your legs.

Your operative leg (leg you had surgery on) turns as a whole unit. Your foot, knee and hip should turn together (see Figure 9).




Figure 9. Sleeping positions after a hip replacement

Do not sleep on your back or unoperated side without pillows between your legs. Do not lie on your operated side (the side you had surgery on) until your doctor tells you it's OK.

Exercising Your Lungs

It's important to exercise your lungs so they expand fully. This helps prevent pneumonia.

- Use your incentive spirometer 10 times every hour you're awake. For more information, read *How to Use Your Incentive Spirometer*. You can find it in the "Educational Resources" section of this guide.
- Do coughing and deep breathing exercises. A member of your care team will teach you how.

Eating and Drinking

Eating a balanced diet high in protein will help you heal after surgery. Your diet should include a healthy protein source at each meal, as well as fruits, vegetables, and whole grains.

For more tips on increasing the amount of calories and protein in your diet, read our resource *Eating Well During Your Cancer Treatment*.

If you have questions about your diet, ask to see a clinical dietitian nutritionist.

Caring for Your Tubes and Drains

Your drainage tubes usually remain in place for a few days after surgery. How long you have them in for depends on how much drainage you're having. Your nurse will measure your drainage 3 times a day. Your doctor will decide when the amount is small enough to remove the drain.

Removing the drainage tubes is usually a quick procedure. It's done in your room and you won't need pain medication. You may feel a stinging sensation, which lasts only a few minutes.

Planning for Discharge

You'll have a case manager who will help you with your discharge planning. Some people go to a short-term rehabilitation center after they leave the hospital. Others can go right home. Everyone's plan is different. Your care team, PT, OT, and case manager will work with you to choose the plan best for you. Your case manager can make arrangements for any equipment, nursing, or rehabilitation you'll need after you leave the hospital.

You may also have an ultrasound done before you leave the hospital. This will help us see if you have a blood clot. We can then decide what blood thinners you should take when you go home. You'll also get antibiotics in your IV to prevent infection.

Your first appointment will be about 2 weeks after you leave the hospital. Call your healthcare provider's office as soon as possible after you leave the hospital to make the appointment.

Leaving the Hospital

Before you leave, look at your incision with one of your healthcare providers. Knowing what it looks like will help you notice any changes later.

On the day of your discharge, plan to leave the hospital around 11 a.m. Before you leave, your healthcare provider will write your discharge order and prescriptions. You'll also get written discharge instructions. One of your healthcare providers will review them with you before you leave.

If your ride is not at the hospital when you're ready to leave, you may be able to wait in the Patient Transition Lounge. A member of your care team will give you more information.

At Home

Read *What You Can Do to Avoid Falling* to learn what you can do to keep from falling at home and during your appointments at MSK. You can find it at www.mskcc.org/pe/avoid_falling or ask your healthcare provider for a copy.

Managing Your Pain

People have pain or discomfort for different lengths of time. You may still have some pain when you go home and will probably be taking pain medication. Some people have soreness, tightness, or muscle aches around their incision for 6 months or longer. This does not mean something is wrong.

Follow these guidelines to help manage your pain at home.

- Take your medications as directed and as needed.
- Call your healthcare provider if the medication prescribed for you does not help your pain.

- Do not drive or drink alcohol while you're taking prescription pain medication. Some prescription pain medications can make you drowsy (very sleepy). Alcohol can make the drowsiness worse.
- As your incision heals, you'll have less pain and need less pain medication. An over-the-counter pain reliever will help with aches and discomfort. Acetaminophen (Tylenol®) and ibuprofen (Advil® or Motrin®) are examples of over-the-counter pain relievers.
 - Follow your healthcare provider's instructions for stopping your prescription pain medication.
 - Do not take too much of any medication. Follow the instructions on the label or from your healthcare provider.
 - Read the labels on all the medications you're taking. This is very important if you're taking acetaminophen. Acetaminophen is an ingredient in many over-the-counter and prescription medications. Taking too much can harm your liver. Do not take more than one medication that has acetaminophen without talking with a member of your care team.
- Pain medication should help you get back to your normal activities. Take enough medication to do your activities and exercises comfortably. It's normal for your pain to increase a little as you start to be more active.
- Keep track of when you take your pain medication. It works best 30 to 45 minutes after you take it. Taking it when you first have pain is better than waiting for the pain to get worse.

Some prescription pain medications (such as opioids) may cause constipation. Constipation is when you poop less often than usual, have a harder time pooping, or both.

Preventing and Managing Constipation

Talk with your healthcare provider about how to prevent and manage constipation. You can also follow these guidelines.

- Go to the bathroom at the same time every day. Your body will get used to going at that time. But if you feel like you need to go, don't put it off.
- Try to use the bathroom 5 to 15 minutes after meals. After breakfast is a good time to go. That's when the reflexes in your colon are strongest.
- Exercise, if you can. Walking is an excellent form of exercise.
- Drink 8 to 10 (8-ounce) cups (2 liters) of liquids daily, if you can. Choose water, juices (such as prune juice), soups, and milkshakes. Avoid liquids with caffeine, such as coffee and soda. Caffeine can pull fluid out of your body.
- Slowly increase the fiber in your diet to 25 to 35 grams per day. Unpeeled fruits and vegetables, whole grains, and cereals contain fiber. If you have an ostomy or recently had bowel surgery, ask your healthcare provider before changing your diet.
- Both over-the-counter and prescription medications can treat constipation. Ask your healthcare provider before taking any medications for constipation. This is very important if you have an ostomy or have had bowel surgery. Follow the instructions on the label or from your healthcare provider. Examples of over-the-counter medications for constipation are:
 - Docusate sodium (Colace®). This is a stool softener (medication that makes your bowel movements softer) that causes few side effects. You can use it to help prevent constipation. Do not take it with mineral oil.
 - Polyethylene glycol (MiraLAX®). This is a laxative (medication that causes bowel movements) that causes few side effects. Take it with 8 ounces (1 cup) of a liquid. Only take it if you're already constipated.
 - Senna (Senokot®). This is a stimulant laxative, which can cause cramping. It's best to take it at bedtime. Only take it if you're already constipated.

If any of these medications cause diarrhea (loose, watery bowel movements), stop taking them. You can start again if needed.

Caring for Your Incision

Before you leave the hospital, you will get information from your healthcare provider. It will tell you how to take care of your dressing when you are home. You may be told to change your dressing every day or 1 week after surgery. You may also be told to leave the dressing on until your surgeon sees you at your post-op (after surgery) appointment.

This is usually 2 to 3 weeks after your surgery. Do not put lotions or creams on your incision unless your healthcare provider tells you to.

It's common for the skin below your incision to feel numb. This happens because some of your nerves were cut during your surgery. The numbness will go away over time.

Call your healthcare provider's office if:

- The skin around your incision is very red or getting more red.
- The skin around your incision is warmer than usual.
- The area around your incision is starting to swell or getting more swollen.
- You see drainage that looks like pus (thick and milky).
- Your incision smells bad.

If you go home with staples in your incision, your healthcare provider will take them out during your first appointment after surgery.

If you have Steri-Strips or Dermabond on your incision, they'll loosen and fall or peel off on their own. If they haven't fallen off after 10 days, you can take them off.

Showering

Do not get your incision and dressing wet until you see your healthcare provider at your post-op appointment. You can sponge bathe. Keep your dressing dry by covering it with a "cast bag," or with tape and a clean garbage bag or plastic wrap. Your healthcare provider will tell you when you can start getting your incision wet.

Eating and Drinking

You can eat all the foods you did before your surgery, unless your healthcare provider gives you other instructions. Eating a balanced diet with lots of calories and protein will help you heal after surgery. Try to eat a good protein source (such as meat, fish, or eggs) at each meal. You should also try to eat fruits, vegetables, and whole grains.

It's also important to drink plenty of liquids. Choose liquids without alcohol or caffeine. Try to drink 8 to 10 (8-ounce) cups of liquids every day.

If you have questions about your diet, ask to see a clinical dietitian nutritionist.

Physical Activity and Exercise

When you leave the hospital, your incision may look like it's healed on the outside. It will not be healed on the inside. Do the following unless your healthcare provider says it's OK.

- Do not lift anything heavier than 10 pounds (4.5 kilograms).
- Do not do any high-energy activities, such as jogging and tennis.
- Do not play any contact sports, such as football.

After your surgery, you must follow special hip precautions. If you get more PT or OT after you leave the hospital, your therapists will continue to teach you ways to move safely. This will help you get stronger and comfortable walking and doing your normal activities.

Doing aerobic exercise, such as walking and stair climbing, will help you gain strength and feel better. Walk at least 2 to 3 times a day for 20 to 30 minutes. You can walk outside or indoors at your local mall or shopping center.

It's common to have less energy than usual after surgery. Recovery time is different for each person. You may need a nap during the day, but try to stay out of bed as much as possible. That will help you sleep at night. It usually takes 6 to 8 weeks until your energy level returns to normal. Increase your activities each day as much as you can. Always balance activity periods with rest periods. Rest is an important part of your recovery.

Sexual Activity

Ask your healthcare provider how soon after going home you may resume sexual activity. For most people, it will be 2 weeks after you leave the hospital.

In order to follow hip precautions, the only limitation is on the positions you may use during sexual activity. You may be on the bottom or lie on your side. Keep 2 pillows between your knees to keep a space between them of 6 to 10 inches (15 to 25 centimeters).

Try to plan the timing of your sexual activity for when you're the most comfortable. Some people find they feel better in the morning. Others feel stiff and sore at that time. Taking your pain medication about 45 minutes before the desired time may also be helpful.

Preventing Infections

If you're going to have any procedure that might cause bleeding, you must tell your doctor or dentist that you have a hip prosthesis. You should take an antibiotic. If you get an infection in your body, it may infect your new hip joint. For more information, read our resource *Preventing an Infection in Your Bone or Joint Replacement Prosthesis*. You can or ask your healthcare provider for a copy or find it at www.mskcc.org/cancer-care/pe/preventing-infection-prosthesis.

Driving

Ask your healthcare provider when you can drive. Most people can start driving again 3 months after surgery. Do not drive while you're taking pain medication that may make you drowsy.

You can ride in a car as a passenger at any time after you leave the hospital.

Going Back to Work

Talk with your healthcare provider about your job. They'll tell you when it may be safe for you to start working again based on what you do. If you move around a lot or lift heavy objects, you may need to stay out a little longer. If you sit at a desk, you may be able to go back sooner.

Managing Your Feelings

After surgery for a serious illness, you may have new and upsetting feelings. Many people say they felt weepy, sad, worried, nervous, irritable, and angry at one time or another. You may find that you cannot control some of these feelings.

If this happens, it's a good idea to seek emotional support. Your healthcare provider can refer you to MSK's Counseling Center. You can also reach them by calling 646-888-0200.

The first step in coping is to talk about how you feel. Family and friends can help. Your healthcare providers can reassure, support, and guide you. It's always a good idea to let us know how you, your family, and your friends are feeling emotionally. Many resources are available to you and your family. Whether you're in the hospital or at home, we're here to help you and your family and friends handle the emotional aspects of your illness.

Notes _____

When To Call Your Healthcare Provider



Call your healthcare provider if:

- You have a fever of 100.5 °F (38 °C) or higher.
- You have chills.
- You're having trouble breathing.
- The skin around your incision is very red or getting more red.
- The skin around your incision is warmer than usual.
- The area around your incision is starting to swell or getting more swollen.
- You see drainage that looks like pus (thick and milky).
- Your incision smells bad.
- You have numbness, tingling (pins and needles), or burning of your toes.
- You have trouble moving your toes.
- You have pain that doesn't go away with medication or pain that's getting worse.
- Your toes are very cold and not warmed by covering them.
- You have trouble breathing.
- You have any questions or concerns.

Contact Information

Monday through Friday from 9 a.m. to 5 p.m., call your healthcare provider's office.

After 5 p.m., during the weekend, and on holidays, call 212-639-2000. Ask to speak to the person on call for your healthcare provider.

Support Services

This section has a list of support services. They may help you as you get ready for your surgery and recover after your surgery.

As you read this section, write down questions to ask your healthcare provider. You can use the space below.

Notes _____

MSK Support Services

Visit the Cancer Types section of MSK's website (www.mskcc.org/cancer-care/types) for more information.

Admitting Office

212-639-7606

Call if you have questions about your hospital admission, such as asking for a private room.

Anesthesia

212-639-6840

Call if you have questions about anesthesia.

Blood Donor Room

212-639-7643

Call for information if you're interested in donating blood or platelets.

Bobst International Center

888-675-7722

We welcome patients from around the world and offer many services to help. If you're an international patient, call for help arranging your care.

Counseling Center

646-888-0200

Many people find that counseling helps them. Our counseling center offers counseling for individuals, couples, families, and groups. We can also prescribe medications to help if you feel anxious or depressed. To make an appointment, ask your healthcare provider for a referral or call the number above.

Female Sexual Medicine and Women's Health Program

646-888-5076

Cancer and cancer treatments can affect your sexual health, fertility, or both. Our Female Sexual Medicine and Women's Health Program can help with sexual health problems, such as premature menopause or fertility issues. We can help before, during, or after your treatment. Call for more information or to make an appointment.

Food Pantry Program

646-888-8055

We give food to people in need during their cancer treatment. For more information, talk with your healthcare provider or call the number above.

Integrative Medicine Service

www.mskcc.org/integrativemedicine

Our Integrative Medicine Service offers many services to complement (go along with) traditional medical care, including music therapy, mind/body therapies, dance and movement therapy, yoga, and touch therapy. To schedule an appointment for these services, call 646-449-1010.

You can also schedule a consultation with a healthcare provider in the Integrative Medicine Service. They will work with you to come up with a plan for creating a healthy lifestyle and managing side effects. To make an appointment, call 646-608-8550.

Male Sexual and Reproductive Medicine Program

646-888-6024

Cancer and cancer treatments can affect your sexual health, fertility, or both. Our Male Sexual and Reproductive Medicine Program can help with sexual health problems, such as erectile dysfunction (ED). We can help before, during, or after your treatment. Call for more information or to make an appointment.

MSK Library

library.mskcc.org

212-639-7439

You can visit our library website or call to talk with the library reference staff. They can help you find more information about a type of cancer. You can also visit LibGuides on MSK's library website at libguides.mskcc.org

Nutrition Services

www.mskcc.org/nutrition

212-639-7312

Our Nutrition Service offers nutritional counseling with one of our clinical dietitian nutritionists. Your clinical dietitian nutritionist will talk with you about your eating habits. They can also give advice on what to eat during and after treatment. To make an appointment, ask a member of your care team for a referral or call the number above.

Patient and Caregiver Education

www.mskcc.org/pe

Visit our patient and caregiver education website to search for educational resources, videos, and online programs.

Patient and Caregiver Peer Support Program

212-639-5007

It can be comforting to talk with someone who has been through a treatment like yours. You can talk with a former MSK patient or caregiver through our Patient and Caregiver Peer Support Program. Your conversations are private. They can be in person or over the phone.

Patient Billing

646-227-3378

Call if you have questions about preauthorization with your insurance company. This is also called preapproval.

Patient Representative Office

212-639-7202

Call if you have questions about the Health Care Proxy form or concerns about your care.

Perioperative Nurse Liaison

212-639-5935

Call if you have questions about MSK releasing any information while you're having surgery.

Private Duty Nurses and Companions

917-862-6373

You can request private nurses or companions to care for you in the hospital and at home. Call for more information.

Resources for Life After Cancer (RLAC) Program

646-888-8106

At MSK, care does not end after your treatment. The RLAC Program is for patients and their families who have finished treatment.

This program has many services. We offer seminars, workshops, support groups, and counseling on life after treatment. We can also help with insurance and employment issues.

Social Work

www.mskcc.org/socialwork

212-639-7020

Social workers help patients, families, and friends deal with common issues for people who have cancer. They provide individual counseling and support groups throughout your treatment. They can help you communicate with children and other family members.

Our social workers can also help refer you to community agencies and programs. They also have information about financial resources, if you're having trouble paying your bills. Call the number above for more information.

Spiritual Care

212-639-5982

Our chaplains (spiritual counselors) are available to listen, help support family members, and pray. They can contact community clergy or faith groups, or simply be a comforting companion and a spiritual presence. Anyone can ask for spiritual support. You do not have to have a religious affiliation (connection to a religion).

MSK's interfaith chapel is located near Memorial Hospital's main lobby. It's open 24 hours a day. If you have an emergency, call 212-639-2000. Ask for the chaplain on call.

Tobacco Treatment Program

www.mskcc.org/tobacco

212-610-0507

If you want to quit smoking, MSK has specialists who can help. Call for information.

Virtual Programs

www.mskcc.org/vp

We offer online education and support for patients and caregivers. These are live sessions where you can talk or just listen. You can learn about your diagnosis, what to expect during treatment, and how to prepare for your cancer care.

Sessions are private, free, and led by experts. Visit our website for more information about Virtual Programs or to register.

External Support Services

There are many other support services available to help you before, during, and after your cancer treatment. Some offer support groups and information. Others can help with transportation, lodging, and treatment costs.

Visit www.mskcc.org/pe/external_support_services for a list of these support services. You can also talk with an MSK social worker by calling 212-639-7020.

Notes _____

Educational Resources

This section has the educational resources mentioned in this guide. They will help you get ready for your surgery and recover after your surgery.

As you read these resources, write down questions to ask your healthcare provider. You can use the space below.

Notes _____



PATIENT & CAREGIVER EDUCATION

How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil

This information will help you check if your medicines or dietary supplements have aspirin, other NSAIDs, vitamin E, or fish oil as an active ingredient. NSAID stands for nonsteroidal anti-inflammatory drug.

It's important to stop taking these medicines and supplements before many cancer treatments. They affect your platelets (blood cells that clot to prevent bleeding) and can raise your risk of bleeding.

Other dietary supplements, such as vitamins and herbal remedies, can also affect your cancer treatment. Read *Herbal Remedies and Cancer Treatment* (www.mskcc.org/pe/herbal_remedies) to learn more.

Make sure your healthcare provider always knows all the prescription and over-the-counter medicines and supplements you're taking. This includes patches and creams.

A prescription medicine is one you can only get with a prescription from your healthcare provider. An over-the-counter medicine is one you can buy without a prescription.

What is an active ingredient?

An active ingredient is the part of a medicine or supplement that makes it work. Some medicines and supplements have just one active ingredient. Others have more. For example:

- Ibuprofen is the active ingredient in Advil® and Motrin®. Ibuprofen is an NSAID.
- Naproxen is the active ingredient in Aleve®. Naproxen is an NSAID.
- Acetaminophen is the active ingredient in Tylenol®.
- Aspirin, acetaminophen, and caffeine are the active ingredients in Excedrin®.

Generic medicines sometimes use their active ingredient as their name. But people often call medicines and supplements by a brand name, even if they're generic. This can make it hard to know their active ingredients.

How to find a medicine or supplement's active ingredients

You can always find the active ingredients by reading the label.

Over-the-counter medicines

Over-the-counter medicines list their active ingredients in the “Drug Facts” label (see Figure 1). Active ingredients are always the first thing on the Drug Facts label.




Figure 1. Active ingredients on an over-the-counter medicine label

Prescription medicines

Prescription medicines list their active ingredients on the label. Their active ingredients and their generic name are the same thing.

Labels often look different depending on which pharmacy you use. Here's an example of where to find a medicine's active ingredients (generic name) on a label from MSK's pharmacy (see Figure 2).




Figure 2. Active ingredients on a prescription medicine label

Dietary supplements

Dietary supplements list their active ingredients in the "Supplement Facts" label (see Figure 3). The active ingredients always have an amount per serving and % daily value included.

	Amount Per Serving	% Daily Value
Vitamin A (as retinyl acetate and 50% as beta-carotene)	5000 IU	100%
Vitamin C (as ascorbic acid)	60 mg	100%
Vitamin D (as cholecalciferol)	400 IU	100%
Vitamin E (as di-alpha tocopheryl acetate)	30 IU	100%
Thiamin (as thiamin mononitrate)	1.5 mg	100%
Riboflavin	1.7 mg	100%
Niacin (as niacinamide)	20 mg	100%
Vitamin B ₆ (as pyridoxine hydrochloride)	2.0 mg	100%
Folate (as folic acid)	400 mcg	100%
Vitamin B ₁₂ (as cyanocobalamin)	6 mcg	100%
Biotin	30 mcg	10%
Pantothenic Acid (as calcium pantothenate)	10 mg	100%

Other ingredients: Gelatin, lactose, magnesium stearate, microcrystalline cellulose, FD&C Yellow No. 6, propylene glycol, propylparaben, and sodium benzoate.

Figure 3. Active ingredients on a supplement label

Active ingredients to look for

If your medicine or supplement has any of these active ingredients, you may need to stop taking it before, during, or after your cancer treatment or surgery. Follow your care team's instructions.

Active ingredients to look for		
<ul style="list-style-type: none">• Acetylsalicylic acid• Alpha-linolenic acid (ALA)• Aspirin• Acetaminophen*• Celecoxib• Diclofenac• Diflunisal• Docosahexaenoic acid (DHA)• Eicosapentaenoic acid (EPA)	<ul style="list-style-type: none">• Etodolac• Fish oil• Fenoprofen Flurbiprofen• Ibuprofen• Indomethacin• Ketoprofen• Ketorolac• Meclofenamate• Mefenamic acid• Meloxicam	<ul style="list-style-type: none">• Nabumetone• Naproxen• Omega-3 fatty acids• Omega-6 fatty acids• Oxaprozin• Piroxicam• Sulindac• Tolmetin• Vitamin E

* The full name acetaminophen isn't always written out. Look for the common abbreviations listed below, especially on prescription pain relievers.

Common abbreviations for acetaminophen		
<ul style="list-style-type: none">• APAP• Acetamin	<ul style="list-style-type: none">• AC• Acetam	<ul style="list-style-type: none">• Acetaminop• Acetaminoph

About acetaminophen (Tylenol)

In general, acetaminophen is safe to take during cancer treatment. It doesn't affect platelets. That means it will not raise your chance of bleeding. If you're getting chemotherapy, talk with your healthcare provider before taking acetaminophen.

There is a limit to how much acetaminophen you can take in a day. Always follow the instructions from your care team or on the medicine's label.

Acetaminophen is in many different prescription and over-the-counter medicines. It's possible to take too much without knowing. **Always read the label on the medicines you take.** Do not take more than 1 medicine that has acetaminophen at a time without talking with a member of your care team.

Instructions before your cancer treatment

Tell your healthcare provider if you take aspirin, other NSAIDs, vitamin E, or fish oil. They'll tell you if you need to stop taking it. You'll also find instructions in the information about your treatment.

Before your surgery

Follow these instructions if you're having surgery or a surgical procedure. **If your healthcare provider gives you other instructions, follow those instead.**

- If you take aspirin or a medicine that has aspirin, you may need to change your dose or stop taking it 7 days before your surgery. Follow your healthcare provider's instructions. **Do not stop taking aspirin unless your healthcare provider tells you to.**
- If you take vitamin E, fish oil, or a supplement that has vitamin E or fish oil, stop taking it 7 days before your surgery or as directed by your healthcare provider.
- If you take an NSAID or a medicine that has an NSAID, stop taking it 48 hours (2 days) before your surgery or as directed by your healthcare provider.

Before your radiology procedure

Follow these instructions if you're having a radiology procedure (including Interventional Radiology, Interventional Mammography, Breast Imaging, and General Radiology). **If your healthcare provider gives you other instructions, follow those instead.**

- If you take aspirin or a medicine that has aspirin, you may need to stop taking it 5 days before your procedure. Follow your healthcare provider's instructions. **Do not stop taking aspirin unless your healthcare provider tells you to.**
- If you take an NSAID or a medicine that has an NSAID, you may need to stop taking it 24 hours (1 day) before your procedure. Follow your healthcare provider's instructions.

Before and during your chemotherapy

Chemotherapy can lower your platelet count, which can increase your risk of bleeding. No matter if you're just starting chemotherapy or have been getting it, talk with your healthcare provider before taking aspirin, other NSAIDs, vitamin E, or fish oil.

If you have any questions, contact a member of your care team directly. If you're a patient at MSK and you need to reach a provider after 5 p.m., during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

How To Check if a Medicine or Supplement Has Aspirin, Other NSAIDs, Vitamin E, or Fish Oil - Last updated on November 29, 2023

All rights owned and reserved by Memorial Sloan Kettering Cancer Center

PATIENT & CAREGIVER EDUCATION

Herbal Remedies and Cancer Treatment

This information explains herbal remedies and how they can affect your treatment.

About Herbal Remedies

Herbal remedies are any herbs, botanical (plant-based) supplements, or dietary supplements you take for their health benefits. These may come as tablets, capsules, powders, teas, liquid extracts, and fresh or dried plants.

Some herbal remedies can help prevent or manage side effects of cancer or your treatment. The herbal remedies that can help you depend on what symptoms you have and what treatment you're getting.

Even though herbal remedies can feel safe, they may not all be safe. Herbal remedies do not go through the same testing as prescription medications to make sure they work and are safe.

Some herbal remedies may be harmful. This is because they can:

- Affect how your other medications work.
- Raise or lower your blood pressure.
- Thin your blood and increase your risk of bleeding.
- Keep radiation therapy from working as well as it should.
- Change how your body reacts to sedation (medication to make you calmer) or general anesthesia (medication to make you sleepy).

Talk with your healthcare provider about any herbal remedies or other

supplements you are taking. They can provide an open and safe space to talk about these products.

For more information about herbs and supplements, visit www.aboutherbbs.com or call MSK's Integrative Medicine Service at 646-608-8550.

Stop taking herbal remedies before your treatment

Stop taking herbal remedies and other dietary supplements 7 days (1 week) before you:

- Have surgery.
- Start chemotherapy.
- Start radiation therapy.
- Have certain procedures. Your healthcare provider will let you know if you need to stop taking herbal remedies before your procedure.

Herbal remedies and other dietary supplements can cause bleeding and affect your treatment. Follow your healthcare provider's instructions for when to restart taking herbal remedies.

You can still use some herbs in your food and drinks, such as using spices in cooking and drinking tea. Herbal remedies are stronger than the herbs you cook with.

Common Herbal Remedies and Their Effects

These are some commonly used herbs and their side effects on cancer treatments.

Echinacea (EH-kih-NAY-shuh)

- Can cause rare but serious allergic reactions, such as a rash or trouble breathing.
- Can keep medications that weaken your immune system from working as well as they should.

Garlic

- Can lower your blood pressure and cholesterol levels.
- Can increase your risk of bleeding.

Ginkgo (also known as Ginkgo biloba)

- Can increase your risk of bleeding.

Ginseng (JIN-seng)

- Can keep sedation or general anesthesia from working as well as they should.
- Can increase your blood pressure.
- Can increase your risk of bleeding.
- Can lower your blood glucose (sugar) level.

Turmeric (TER-mayr-ik)

- Can keep chemotherapy from working as well as it should.

St. John's Wort

- Can keep some medications from working as well as they should.
- Can make your skin more sensitive to radiation or laser treatment.

Valerian (vuh-LEER-ee-un)

- Can make sedation or general anesthesia affect you more than they should.

Herbal formulas

- Herbal formulas contain many different herbs and dosages.
- Stop taking these products 7 days (1 week) before treatment. Do not start taking herbal formulas again until your healthcare provider tells you it is safe.

This information does not cover all herbal remedies or possible side effects. Talk with your healthcare provider if you have any questions or concerns.

Contact Information

- To schedule a consultation with a healthcare provider in Integrative Medicine, call 646-608-8550.
- To make an appointment for Integrative Medicine Service's therapies, classes, and workshops, call 646-449-1010.

For more information, visit www.mskcc.org/IntegrativeMedicine or read *Integrative Medicine Therapies and Your Cancer Treatment* (www.mskcc.org/pe/integrative_therapies).

For more resources, visit www.mskcc.org/pe to search our virtual library.

Herbal Remedies and Cancer Treatment - Last updated on May 5, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



PATIENT & CAREGIVER EDUCATION

How To Use Your Incentive Spirometer

This information will help you learn how to use your incentive spirometer (in-SEN-tiv spy-rah-MEE-ter). It also answers some common questions about it.

About your incentive spirometer

After your surgery you may feel weak and sore, and it may be uncomfortable to take deep breaths. Your healthcare provider may recommend using a device called an incentive spirometer (see Figure 1). It helps you practice taking deep breaths.




Figure 1. Parts of an incentive spirometer

It's important to use your incentive spirometer after your surgery. Using an incentive spirometer:

- Helps your lungs expand so you can take deep, full breaths.
- Exercises your lungs and makes them stronger as you heal from surgery.

If you have a respiratory infection, do not use your incentive spirometer around other people. A respiratory infection is an infection in your nose, throat, or lungs, such as pneumonia (noo-MOH-nyuh) or COVID-19. This kind of infection can spread from person to person through the air.

How to use your incentive spirometer

Here is a video that shows how to use your incentive spirometer:



Please visit www.mskcc.org/pe/incentive_spirometer_video to watch this video.

Setting up your incentive spirometer

Before you use your incentive spirometer for the first time, you will need to set it up. First, take the flexible (bendable) tubing out of the bag and stretch it out. Then, connect the tubing to the outlet on the right side of the base (see Figure 1). The mouthpiece is attached to the other end of the tubing.

Knowing what number to aim for on your incentive spirometer

Your healthcare provider will teach you how to use your incentive spirometer before you leave the hospital. They will help you set a goal and tell you what number to aim for when using your spirometer. If a goal was not set for you, talk with your healthcare provider. Ask them what number you should aim for.

You can also check the package your incentive spirometer came in. It may have a chart to help you figure out what number to aim for. To learn more, read “What number I should aim for?” in the “Common questions about your

incentive spirometer” section.

Using your incentive spirometer

When using your incentive spirometer, make sure to breathe through your mouth. If you breathe through your nose, your spirometer will not work right.

Follow these steps to use your incentive spirometer. Repeat these steps every hour you’re awake. Follow the instructions from your healthcare provider if they’re different from the ones here.

1. Sit upright in a chair or in bed. Hold your incentive spirometer at eye level.
2. Put the mouthpiece in your mouth and close your lips tightly around it. Make sure you do not block the mouthpiece with your tongue.
3. With the mouthpiece in your mouth, breathe out (exhale) slowly and fully.
 - Some people may have trouble exhaling with the mouthpiece in their mouth. If you do, take the mouthpiece out of your mouth, and then exhale slowly and fully. After you exhale, put the mouthpiece back in your mouth and go on to step 4.
4. Breathe in (inhale) slowly through your mouth, as deeply as you can. You will see the piston slowly rise inside the spirometer. The deeper you breathe in, the higher the piston will rise.
5. As the piston rises, the coaching indicator on the right side of the spirometer should also rise. It should float between the 2 arrows (see Figure 1).
 - The coaching indicator measures the speed of your breath. If it does not stay between the 2 arrows, you’re breathing in either too fast or too slow.
 - If the indicator rises above the higher arrow, you’re breathing in too fast. Try to breathe in slower.
 - If the indicator stays below the lower arrow, you’re breathing in too slow. Try to breathe in faster.

6. When you cannot breathe in any further, hold your breath for at least 3 to 5 seconds. Hold it for longer if you can. You will see the piston slowly fall to the bottom of the spirometer.
7. Once the piston reaches the bottom of the spirometer, breathe out slowly and fully through your mouth. If you want, you can take the mouthpiece out of your mouth first and then breathe out.
8. Rest for a few seconds. If you took the mouthpiece out of your mouth, put it back in when you're ready to start again.
9. Repeat steps 1 to 8 at least 10 times. Try to get the piston to the same level with each breath. After you have done the exercise 10 times, go on to step 10.
10. Use the marker on the left side of the spirometer to mark how high the piston rises (see Figure 1). **Look at the very top of the piston, not the bottom. The number you see at the top is the highest number the piston reached. Put the marker there.** This is how high you should try to get the piston the next time you use your spirometer.
 - Write down the highest number the piston reached. This can help you change your goals and track your progress over time.

Take 10 breaths with your incentive spirometer every hour you're awake.

Cover the mouthpiece of your incentive spirometer when you're not using it.

Tips for using your incentive spirometer

Follow these tips when using your incentive spirometer:

- If you had surgery on your chest or abdomen (belly), it may help to splint your incision (surgical cut). To do this, hold a pillow firmly against your incision. This will keep your muscles from moving as much while you're using your incentive spirometer. It will also help ease pain at your incision.
- If you need to clear your lungs, you can try to cough a few times. As

you're coughing, hold a pillow against your incision, as needed.

- If you feel dizzy or lightheaded, take the mouthpiece out of your mouth. Then, take a few normal breaths. Stop and rest for a while, if needed. When you feel better, you can go back to using your incentive spirometer.
- You may find it hard to use your incentive spirometer at first. If you cannot make the piston rise to the number your healthcare provider set for you, it's OK. Reaching your goal takes time and practice. It's important to keep using your spirometer as you heal from surgery. The more you practice, the stronger your lungs will get.

Common questions about your incentive spirometer

How often should I use my incentive spirometer?

How often you will need to use your incentive spirometer is not the same for everyone. It depends on the type of surgery you had and your recovery process.

Most people can take 10 breaths with their spirometer every hour they're awake.

Your healthcare provider will tell you how often to use your spirometer. Follow their instructions.

How long after my surgery will I need to use my incentive spirometer?

The length of time you will need to use your incentive spirometer is not the same for everyone. It depends on the type of surgery you had and your recovery process.

Your healthcare provider will tell you how long you need to use your spirometer. Follow their instructions.

How do I clean my incentive spirometer?

An incentive spirometer is a disposable device and only meant to be used for a short time. Because of this, you may not find cleaning instructions in the package your spirometer came in. If you have questions about cleaning your spirometer, talk with your healthcare provider.

What do the numbers on my incentive spirometer measure?

The large column of your incentive spirometer has numbers on it (see Figure 1). These numbers measure the volume of your breath in milliliters (mL) or cubic centimeters (cc). The volume of your breath is how much air you can breathe into your lungs (inhale).

For example, if the piston rises to 1500, it means you can inhale 1500 mL or cc of air. The higher the number, the more air you're able to inhale, and the better your lungs are working.

What number I should aim for?

The number you should aim for depends on your age, height, and sex. It also depends on the type of surgery you had and your recovery process. Your healthcare provider will look at these things when setting a goal for you. They will tell you what number to aim for.

Most people start with a goal of 500 mL or cc. Your healthcare provider may change your goal and have you aim for higher numbers as you heal from surgery.

The package your incentive spirometer came in may have a chart. You can use the chart to set your goal based on your age, height, and sex. If you cannot find this information, ask your healthcare provider what your goal should be.

What does the coaching indicator on my incentive spirometer measure?

The coaching indicator on your incentive spirometer measures the speed of your breath. As the speed of your breath changes, the indicator moves up and down.

Use the indicator to guide your breathing. If the indicator rises above the higher arrow, it means you're breathing in too fast. If the indicator stays below the lower arrow, it means you're breathing in too slow.

Aim to keep the indicator between the 2 arrows (see Figure 1). This means your breath is steady and controlled.

When to call your healthcare provider

Call your healthcare provider if you have any of these when using your incentive spirometer:

- Feel dizzy or lightheaded.
- Pain in your lungs or chest.
- Severe (very bad) pain when you take deep breaths.
- Trouble breathing.
- Coughing up blood.
- Fluid or blood coming from your incision site when you cough.
- Trouble using your spirometer for any reason.

If you have any questions, contact a member of your care team directly. If you're a patient at MSK and you need to reach a provider after 5 p.m., during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

How To Use Your Incentive Spirometer - Last updated on November 24, 2023

All rights owned and reserved by Memorial Sloan Kettering Cancer Center