Ready to start planning your care? Call us at 800-525-2225 to make an appointment.

×



Memorial Sloan Kettering Cancer Center

Make an Appointment Back

Cervical Capper Tratment Refer a Patient

ABOUT US Our mission, vision & core values Leadership History Equality, diversity & inclusion Annual report Give to MSK

time of your surgery that suggest an increased risk of the cancer returning, such as cancer cells in your lymph nodes or in parts of the tissue that is being surgically removed.

When radiation and chemotherapy are recommended in place of surgery, our doctors typically combine external-beam radiation therapy with internal radiation therapy. External-beam radiation therapy consists of x-ray treatments given daily for about five weeks. Low doses of chemotherapy drugs such as cisplatin (Platinol®) are used during this treatment to sensitize the cancer cells to radiation therapy.

Internal radiation is then given using high-dose brachytherapy, in which high doses of radioactive material are applied through a specialized applicator directly to the tumor. Studies have shown that this combined action can dramatically lengthen life in women with advanced cervical cancer. ⁽¹⁾, ⁽²⁾, ⁽³⁾

Specialized approaches to external-beam radiation therapy may include the use of <u>intensity-modulated radiation therapy (IMRT)</u> as well. IMRT is a type of external-beam radiation that enables the doctor to customize the intensity of each radiation beam to conform to specific tumor shapes and sizes. IMRT not only reduces the dosage of radiation to healthy tissues but may also lessen your risk for side effects.

Treatment for Advanced or Recurrent Cancer

If your cancer has spread beyond the pelvis and into organs such as the lungs or liver, or you have recurrent disease, we will work with you to reduce your cancer-related symptoms and help you maintain your quality of life.

We primarily recommend chemotherapy for women with advanced or recurrent disease. Several drugs, such as cisplatin (Platinol®) and paclitaxel (Taxol®), are available. Memorial Sloan Kettering investigators are researching additional chemotherapy strategies that may be effective for patients.

Some women with recurrent cervical cancer choose to undergo an involved surgery known as pelvic exenteration. With this approach, surgeons remove cancerous tissue and reconstruct the remaining organs to preserve as much function as possible. Brachytherapy can be done at the same time. We reserve this radical procedure for women who have otherwise limited treatment options.

Morris M, Eifel PJ, Lu J, et al. Pelvic radiation with concurrent chemotherapy compared with pelvic and para-aortic radiation for high-risk cervical cancer. N Engl J Med. 1999 Apr 15:340(15):1137-43.

Keys HM, Bundy BN, Stehman FB, et al. Cisplatin, radiation, and adjuvant hysterectomy compared with radiation and adjuvant hysterectomy for bulky stage IB cervical carcinoma. N Engl J Med. 1999 Apr 15;340(15):1154-61.

Rose PG, Bundy BN, Watkins EB, et al. Concurrent cisplatin-based radiotherapy and chemotherapy for locally advanced cervical cancer. N

Request an Appointment

Call 800-525-2225 Available Monday through Friday, 8 a.m. to 6 p.m. (Eastern time)

Make an Appointment

PREVIOUS Cervical Cancer Surgery

NEXT

Cervical Cancer Follow-Up Care

Connect
Contact us
Locations
APPOINTMENTS
<u>800-525-2225</u>
About MSK
<u>About us</u>
Cancer Care
Adult cancer types
Child & teen cancer types Integrative medicine
Nutrition & cancer
Find a doctor
Research & Education
Sloan Kettering Institute
Gerstner Sloan Kettering Graduate School

Radiation Therapy & Chemotherapy for Cervical Cancer

MSK Library 💻

Communication preferences Cookie preferences Legal disclaimer Accessibility statement Privacy policy Price transparency Public notices © 2024 Memorial Sloan Kettering Cancer Center