

Ready to start planning your care? Call us at [646-926-0945](tel:646-926-0945) to make an appointment.

×



Memorial Sloan Kettering  
Cancer Center

[Make an Appointment](#)  
[Back](#)

[Prostate Cancer Screening & Treatment](#)  
[Learn About Cancer & Treatment](#)

## ABOUT US

[Our mission, vision & core values](#)

[Leadership](#)

[History](#)

[Inclusion & belonging](#)

[Annual report](#)

[Give to MSK](#)

7 years later with a single measure of prostate-specific antigen at or before age 50. Cancer. Mar 15 2011;117(6):1210-1219.

[Vickers AJ, Till C, Tangen CM, Lilja H, Thompson IM. An empirical evaluation of guidelines on prostate-specific antigen velocity in prostate cancer detection. Journal of the National Cancer Institute. Mar 16 2011;103\(6\):462-469 .](#)

[Roobol MJ, Roobol DW, Schroder FH. Is additional testing necessary in men with prostate-specific antigen levels of 1.0 ng/mL or less in a population-based screening setting? \(ERSPC, section Rotterdam\). Urology. Feb 2005;65\(2\):343-346 .](#)

[Bul M, van Leeuwen PJ, Zhu X, Schroder FH, Roobol MJ. Prostate cancer incidence and disease-specific survival of men with initial prostate-specific antigen less than 3.0 ng/ml who are participating in ERSPC Rotterdam. Eur Urol. Apr 2011;59\(4\):498-505 .](#)

Vickers AJ, Cronin AM, Bjork T, et al. Prostate specific antigen concentration at age 60 and death or metastasis from prostate cancer: case-control study. BMJ. 2010;341:c4521.

## Request an Appointment

Call [800-525-2225](tel:800-525-2225)

We're available 24 hours a day, 7 days a week

[Make an appointment](#)

© 2026 Memorial Sloan Kettering Cancer Center