



Make an Appointment

Prostate Cancer Screening Guidelines Light and Balbeom over 12 Treatment

Refer a Patient

ABOUT US

Our mission, vision & core values

Leadership

History

Equality, diversity & inclusion

Annual report

Give to MSK

single measure of prostate-specific antigen at or before age 50. Cancer. Mar 15 2011;117(6):1210-1219.

<u>Vickers AJ, Till C, Tangen CM, Lilja H, Thompson IM. An empirical evaluation of guidelines on prostate-specific antigen velocity in prostate cancer detection. Journal of the National Cancer Institute. Mar 16 2011;103(6):462-469.</u>

Roobol MJ, Roobol DW, Schroder FH. Is additional testing necessary in men with prostate-specific antigen levels of 1.0 ng/mL or less in a population-based screening setting? (ERSPC, section Rotterdam). Urology. Feb 2005;65(2):343-346.

Bul M, van Leeuwen PJ, Zhu X, Schroder FH, Roobol MJ. Prostate cancer incidence and disease-specific survival of men with initial prostate-specific antigen less than 3.0 ng/ml who are participating in ERSPC Rotterdam. Eur Urol. Apr 2011;59(4):498-505.

Vickers AJ, Cronin AM, Bjork T, et al. Prostate specific antigen concentration at age 60 and death or metastasis from prostate cancer: case-control study. BMJ. 2010;341:c4521.

Request an Appointment

Call 800-525-2225

Available Monday through Friday, 8 a.m. to 6 p.m. (Eastern time)

Make an Appointment

•

- Connect	
Contact us	
<u>Locations</u>	
APPOINTMENTS	
800-525-2225	
- About MSK	
About us	
<u>Careers</u>	
Giving	
Cancer Care	
Adult cancer types	
Child & teen cancer types	
Integrative medicine	
Nutrition & cancer	
Find a doctor	
- Research & Education	
Sloan Kettering Institute	
Gerstner Sloan Kettering Graduate School.■	
Graduate medical education	
MSK Library	
Communication preferences	
Cookie preferences	
Legal disclaimer	
Accessibility statement	
Privacy policy Price transparency	
Public notices	