

Ready to start planning your care? Call us at [800-525-2225](tel:800-525-2225) to make an appointment.

×



Memorial Sloan Kettering
Cancer Center

[Make an Appointment](#)

[Back](#)

[Uterine \(Endometrial\) Cancer](#)

[Refer a Patient](#)

ABOUT US

[Our mission, vision & core values](#)

[Leadership](#)

[History](#)

[Equality, diversity & inclusion](#)

[Annual report](#)

[Give to MSK](#)

are between the ages of 50 and 60

began menstruating before age 12

entered menopause relatively late, after age 52

never gave birth

have a history of infertility (an inability to become pregnant)

have an ovarian disease, such as polycystic ovarian syndrome, that could cause you to have higher than normal levels of the hormone estrogen and lower than normal levels of the hormone progesterone

have elevated blood sugar (diabetes)

have high blood pressure (hypertension)

have a family history of endometrial carcinoma

have taken the drug tamoxifen after menopause; the increased risk depends in part on the dose taken and the length of time it's used. Women who take tamoxifen should discuss the risks and benefits of this drug with their doctors.

have been diagnosed with endometrial hyperplasia

take certain types of hormone replacement therapy (HRT)

Genetic Risk Factors

If several members of your family have had uterine or hereditary nonpolyposis colorectal cancer (HNPCC), also known as [Lynch syndrome](#), consider having genetic counseling and genetic testing through our [Hereditary Cancer & Genetics Service](#).

Women who inherit mutations in the PTEN gene (Cowden's syndrome) may also be at an increased risk of developing endometrial cancer.

Protect Yourself

In addition to getting regular annual checkups with your gynecologist and reporting any unexpected or abnormal vaginal bleeding, lifestyle measures, such as keeping your weight under control with physical activity and following a low-fat diet, may help to prevent uterine cancer. Using oral contraceptives has also been associated with a reduced risk.



Why Black Women Are Twice as Likely to Die of Endometrial Cancer — and What MSK Is Doing to Change It
Learn why Black women face a higher risk of death from endometrial cancer, and about a new effort led by Dr. Carol Brown at Memorial Sloan Kettering Cancer Center to close the gap.

[Learn more](#)

Request an Appointment

Call [800-525-2225](tel:800-525-2225)

Available Monday through Friday, 8 a.m. to 6 p.m. (Eastern time)

[Make an Appointment](#)

PREVIOUS

[Uterine \(Endometrial\) Cancer Signs & Symptoms](#)

NEXT

[Uterine \(Endometrial\) Cancer Diagnosis](#)

▼ Connect

[Contact us](#)

[Locations](#)

APPOINTMENTS

800-525-2225



▼ About MSK

[About us](#)

[Careers](#) ■

[Giving](#) ■

▼ Cancer Care

[Adult cancer types](#)

[Child & teen cancer types](#)

[Integrative medicine](#)

[Nutrition & cancer](#)

[Find a doctor](#)

▼ Research & Education

[Sloan Kettering Institute](#)

[Gerstner Sloan Kettering Graduate School](#) ■

[Graduate medical education](#)

[MSK Library](#) ■

[Communication preferences](#)

[Cookie preferences](#)

[Legal disclaimer](#)

[Accessibility statement](#)

[Privacy policy](#)

[Price transparency](#)

[Public notices](#)

© 2024 Memorial Sloan Kettering Cancer Center