

Ready to start planning your care? Call us at [800-525-2225](tel:800-525-2225) to make an appointment.

×



Memorial Sloan Kettering
Cancer Center

[Make an Appointment](#)

[← Back](#)

[The Prasad Adusumilli Lab](#)

[Research, Diagnosis & Treatment](#)

[Refer a Patient](#)

ABOUT US

[Our mission, vision & core values](#)

[Leadership](#)

[History](#)

[Equality, diversity & inclusion](#)

[Annual report](#)

[Give to MSK](#)

Kagawa University Visiting Research Scholarship

Current Employment

Assistant Professor, Kagawa, University, Japan

Publications

Bains S, Eguchi T, Warth A, Yeh YC, Nitadori JI, Woo KM, Chou TY, Dienemann H, Muley T, Nakajima J, Shinozaki-Ushiku A, Wu YC, Lu S, Kadota K, Jones DR, Travis WD, Tan KS, Adusumilli PS. Procedure-specific risk prediction for recurrence in patients undergoing lobectomy or sublobar resection for small (≤ 2 cm) lung adenocarcinoma: an international cohort analysis. *J Thorac Oncol*. 2019;14(1):72-86.

Kameda K, Eguchi T, Lu S, Qu Y, TanKS, Kadota K, Adusumilli PS, Travis WD. Implications of the eighth edition of the TNM proposal: invasive vs. total tumor size for the T descriptor in pathologic stage I-IIA lung adenocarcinoma. *J Thorac Oncol*. 2018;13(12):1919-1929.

Rosen LE, Karrison T, Ananthanarayanan V, Gallan AJ, Adusumilli PS, Alchami FS, Attanoos R, Brcic L, Butnor KJ, Galateau-Salle F, Hiroshima K, Kadota K, Klampatsa A, Stang NL, Lindenmann J, Litzky LA, Marchevsky A, Medeiros F, Montero MA, Moore DA, Nabeshima K, Pavlisko EN, Roggli VL, Sauter JL, Sharma A, Sheaff M, Travis WD, Vigneswaran WT, Vrugt B, Walts AE, Tjota MY, Krausz T, Husain AN. Nuclear grade and necrosis predict prognosis in malignant epithelioid pleural mesothelioma: a multi-institutional study. *Mod Pathol*. 2018;31(4):598-606.

Bekhtman N, Pietanza CM, Sahari I, Montecalvo J, Wang H, Habeeb O, Kadota K, Adusumilli PS, Rudin CM, Ladanvi

Nonaka H, Hatanaka S, Saito S, Morikawa S, Wang H, Nakao S, Kadota K, Adusumilli PS, Hattori S, Eguchi T, Travis WD, Joubert P. Pulmonary large cell neuroendocrine carcinoma with adenocarcinoma-like features: napsin A expression and genomic alterations. *Mod Pathol*. 2018;31(1):111-21.

Eguchi T, Kadota K, Mayor M, Zauderer MG, Rimmer A, Rusch VW, Travis WD, Sadelain M, Adusumilli PS. Cancer antigen profiling for malignant pleural mesothelioma immunotherapy: expression and coexpression of mesothelin, cancer antigen 125, and Wilms tumor 1. *Oncotarget*. 2017;8(44):77872-82.

Lu S, Tan KS, Kadota K, Eguchi T, Bains S, Rekhtman N, Adusumilli PS, Travis WD. Spread through air spaces (STAS) is an independent predictor of recurrence and lung cancer specific death in squamous cell carcinoma. *J Thorac Oncol*. 2017;12(2):223-234.

Kadota K, Sima CS, Arcila ME, Hedvat C, Kris MG, Jones DR, Adusumilli PS, Travis WD. KRAS mutation is a significant prognostic factor in early-stage lung adenocarcinoma. *Am J Surg Pathol*. 2016;40(12):1579-1590.

Eguchi T, Kadota K, Chaft J, Evans B, Kidd J, See Tan K, Dycoco J, Kolquist K, Davis T, Hamilton SA, Yager K, Jones JT, Travis WD, Jones DR, Hartman AR, Adusumilli PS. Cell cycle progression score is a marker for five-year lung cancer-specific mortality risk in patients with resected stage I lung adenocarcinoma. *Oncotarget*. 2016;7(23):35241-56.

Liu Y, Amin EB, Mayo MW, Chudgar NP, Bucciarelli PR, Kadota K, Adusumilli PS, Jones DR. CK2alpha' drives lung cancer metastasis by targeting BRMS1 nuclear export and degradation. *Cancer Res*. 2016;76(9):2675-86.

Yeh YC, Kadota K, Nitadori J, Sima CS, Rizk NP, Jones DR, Travis WD, Adusumilli PS. International Association for the Study of Lung Cancer/American Thoracic Society/European Respiratory Society classification predicts occult lymph node metastasis in clinically mediastinal node-negative lung adenocarcinoma. *Eur J Cardiothorac Surg*. 2016;49(1):e9-e15.

Kadota K, Eguchi T, Villena-Vargas J, Woo KM, Sima CS, Jones DR, Travis WD, Adusumilli PS. Nuclear estrogen receptor-alpha expression is an independent predictor of recurrence in male patients with pT1aN0 lung adenocarcinomas, and correlates with regulatory T-cell infiltration. *Oncotarget*. 2015;6(29):27505-18.

Ujii H, Kadota K, Chaft JE, Buitrago D, Sima CS, Lee MC, Huang J, Travis WD, Rizk NP, Rudin CM, Jones DR, Adusumilli PS. Solid predominant histologic subtype in resected stage I lung adenocarcinoma is an independent predictor of early, extrathoracic, multisite recurrence and of poor postrecurrence survival. *J Clin Oncol*. 2015;33(26):2877-84.

Kadota K, Nitadori J, Rekhtman N, Jones DR, Adusumilli PS, Travis WD. Reevaluation and reclassification of resected lung carcinomas originally diagnosed as squamous cell carcinoma using immunohistochemical analysis. *Am J Surg Pathol*. 2015;39(9):1170-80.

Kadota K, Villena-Vargas J, Nitadori J, Sima CS, Jones DR, Travis WD, Adusumilli PS. Tumoral CD10 expression correlates with aggressive histology and prognosis in patients with malignant pleural mesothelioma. *Ann Surg Oncol*. 2015;22(9):3136-43.

Kadota K, Yeh YC, Villena-Vargas J, Cherkassky L, Drill EN, Sima CS, Jones DR, Travis WD, Adusumilli PS. Tumor budding correlates with protumor immune microenvironment and is an independent prognostic factor for recurrence of stage I lung adenocarcinoma. *Chest*. 2015;148(3):711-721.

Kadota K, Nitadori J, Ujii H, Buitrago DH, Woo KM, Sima CS, Travis WD, Jones DR, Adusumilli PS. Prognostic impact of immune microenvironment in lung squamous cell carcinoma: tumor-infiltrating CD10+ neutrophil/CD20+ lymphocyte ratio as an independent prognostic factor. *J Thorac Oncol*. 2015;10(9):1301-10.

Kadota K, Buitrago D, Lee MC, Villena-Vargas J, Sima CS, Jones DR, Travis WD, Adusumilli PS. Tumoral CD10 expression correlates with high-grade histology and increases risk of recurrence in patients with stage I lung adenocarcinoma. *Lung Cancer*. 2015;89(3):329-36.

Yeh YC, Nitadori J, Kadota K, Yoshizawa A, Rekhtman N, Moreira AL, Sima CS, Rusch VW, Adusumilli PS, Travis WD. Using frozen section to identify histological patterns in stage I lung adenocarcinoma of ≤ 3 cm: accuracy and interobserver agreement. *Histopathology*. 2015;66(7):922-38.

- Kadota K, Nitadori J, Sima CS, Ujiie H, Rizk NP, Jones DR, Adusumilli PS, Travis WD. Tumor spread through air spaces is an important pattern of invasion and impacts the frequency and location of recurrences after limited resection for small stage I lung adenocarcinomas. *J Thorac Oncol*. 2015;10(5):806-14.
- Buitrago DH, Patnaik SK, Kadota K, Kannisto E, Jones DR, Adusumilli PS. Small RNA sequencing for profiling microRNAs in long-term preserved formalin-fixed and paraffin-embedded non-small cell lung cancer tumor specimens. *PLoS One*. 2015;10(3):e0121521.
- Ujiie H, Kadota K, Nitadori J-i, Aerts JG, Woo KM, Sima CS, Travis WD, Jones DR, Krug LM, Adusumilli PS. Tumoral and stromal immune microenvironment in epithelioid malignant pleural mesothelioma: Tumoral il-7 receptor, tumor-infiltrating cd20+ lymphocytes, and their ratio to tumor-associated macrophages are independent predictors of survival. *Oncoimmunology*. 2015;4(6):e1009285.
- Liu Y, Mayo MW, Xiao A, Hall EH, Amin EB, Kadota K, Adusumilli PS, Jones DR. Loss of BRMS1 promotes a mesenchymal phenotype through NF-kappaB-dependent regulation of Twist1. *Mol Cell Biol*. 2015;35(1):303-17.
- Kadota K, Eguchi T, Villena-Vargas J, Woo KM, Sima CS, Jones DR, Travis WD, Adusumilli PS. Nuclear estrogen receptor-alpha expression is an independent predictor of recurrence in male patients with pT1aN0 lung adenocarcinomas, and correlates with regulatory T-cell infiltration. *Oncotarget*. 2015;6(29):27505-18.
- Lee MC, Buitrago DH, Kadota K, Ujiie H, Woo K, Sima CS, Travis WD, Jones DR, Adusumilli PS. The tumor immune microenvironment in octogenarians with stage I non-small cell lung cancer. *Oncoimmunology*. 2014;3(11):e967142.
- Tozbikian G, Brogi E, Kadota K, Catalano J, Akram M, Patil S, Ho AY, Reis-Filho JS, Weigelt B, Norton L, Adusumilli PS, Wen HY. Mesothelin expression in triple negative breast carcinomas correlates significantly with basal-like phenotype, distant metastases and decreased survival. *PLoS One*. 2014;9(12):e114900.
- Lee MC, Kadota K, Buitrago D, Jones DR, Adusumilli PS. Implementing the new IASLC/ATS/ERS classification of lung adenocarcinomas: results from international and Chinese cohorts. *J Thorac Dis*. 2014;6(Suppl 5):S568-80.
- Eguchi T, Kadota K, Park BJ, Travis WD, Jones DR, Adusumilli PS. The new IASLC-ATS-ERS lung adenocarcinoma classification: what the surgeon should know. *Semin Thorac Cardiovasc Surg*. 2014;26(3):210-22.
- Kadota K, Yeh Y-C, D'Angelo SP, Moreira AL, Kuk D, Sima CS, Riely GJ, Arcila ME, Kris MG, Rusch VW, Adusumilli PS, Travis WD. Associations between mutations and histologic patterns of mucin in lung adenocarcinoma: invasive mucinous pattern and extracellular mucin are associated with KRAS mutation. *Am J Surg Pathol*. 2014;38(8):1118-27.
- Kadota K, Nitadori J, Woo KM, Sima CS, Finley DJ, Rusch VW, Adusumilli PS, Travis WD. Comprehensive pathological analyses in lung squamous cell carcinoma: single cell invasion, nuclear diameter, and tumor budding are independent prognostic factors for worse outcomes. *J Thorac Oncol*. 2014;9(8):1126-39.
- Lee MC, Buitrago DH, Kadota K, Jones DR, Adusumilli PS. Recent advances and clinical implications of the micropapillary histological subtype in lung adenocarcinomas. *Lung cancer management*. 2014;3(3):245-53.
- Kadota K, Yeh YC, Sima CS, Rusch VW, Moreira AL, Adusumilli PS, Travis WD. The cribriform pattern identifies a subset of acinar predominant tumors with poor prognosis in patients with stage I lung adenocarcinoma: a conceptual proposal to classify cribriform predominant tumors as a distinct histologic subtype. *Mod Pathol*. 2014;27(5):690-700.
- Kadota K, Villena-Vargas J, Yoshizawa A, Motoi N, Sima CS, Riely GJ, Rusch VW, Adusumilli PS, Travis WD. Prognostic significance of adenocarcinoma in situ, minimally invasive adenocarcinoma, and nonmucinous lepidic predominant invasive adenocarcinoma of the lung in patients with stage I disease. *Am J Surg Pathol*. 2014;38(4):448-60.
- Kachala SS, Bograd AJ, Villena-Vargas J, Suzuki K, Servais EL, Kadota K, Chou J, Sima CS, Vertes E, Rusch VW, Travis WD, Sadelain M, Adusumilli PS. Mesothelin overexpression is a marker of tumor aggressiveness and is associated with reduced recurrence-free and overall survival in early-stage lung adenocarcinoma. *Clin Cancer Res*. 2014;20(4):1020-8.
- Nitadori J, Bograd AJ, Kadota K, Sima CS, Rizk NP, Morales EA, Rusch VW, Travis WD, Adusumilli PS. Impact of

Nitadori J, Bogdan AG, Kadota K, Sima CS, Rizk NP, Morales EA, Rusch VW, Travis WD, Adusumilli PS. Impact of micropapillary histologic subtype in selecting limited resection vs lobectomy for lung adenocarcinoma of 2cm or smaller. *J Natl Cancer Inst.* 2013;105(16):1212-20.

Kadota K, Nitadori JI, Adusumilli PS. Prognostic value of the immune microenvironment in lung adenocarcinoma. *Oncoimmunology.* 2013;2(5):e24036.

Suzuki K, Kadota K, Sima CS, Nitadori J, Rusch VW, Travis WD, Sadelain M, Adusumilli PS. Clinical impact of immune microenvironment in stage I lung adenocarcinoma: tumor interleukin-12 receptor beta2 (IL-12Rbeta2), IL-7R, and stromal FoxP3/CD3 ratio are independent predictors of recurrence. *J Clin Oncol.* 2013;31(4):490-8.

Kadota K, Nitadori J, Sarkaria IS, Sima CS, Jia X, Yoshizawa A, Rusch VW, Travis WD, Adusumilli PS. Thyroid transcription factor-1 expression is an independent predictor of recurrence and correlates with the IASLC/ATS/ERS histologic classification in patients with stage I lung adenocarcinoma. *Cancer.* 2013;119(5):931-8.

Kadota K, Colovos C, Suzuki K, Rizk NP, Dunphy MP, Zabor EC, Sima CS, Yoshizawa A, Travis WD, Rusch VW, Adusumilli PS. FDG-PET SUVmax combined with IASLC/ATS/ERS histologic classification improves the prognostic stratification of patients with stage I lung adenocarcinoma. *Ann Surg Oncol.* 2012;19(11):3598-605.

Kadota K, Suzuki K, Kachala SS, Zabor EC, Sima CS, Moreira AL, Yoshizawa A, Riely GJ, Rusch VW, Adusumilli PS, Travis WD. A grading system combining architectural features and mitotic count predicts recurrence in stage I lung adenocarcinoma. *Mod Pathol.* 2012;25(8):1117-27.

Kadota K, Kachala SS, Nitadori J, Suzuki K, Dunphy MP, Sima CS, Travis WD, Rusch VW, Adusumilli PS. High SUVmax on FDG-PET indicates pleomorphic subtype in epithelioid malignant pleural mesothelioma: supportive evidence to reclassify pleomorphic as nonepithelioid histology. *J Thorac Oncol.* 2012;7(7):1192-7.

Kadota K, Suzuki K, Colovos C, Sima CS, Rusch VW, Travis WD, Adusumilli PS. A nuclear grading system is a strong predictor of survival in epithelioid diffuse malignant pleural mesothelioma. 2012;25(2):260-71.

Suzuki K, Kachala SS, Kadota K, Shen R, Mo Q, Beer DG, Rusch VW, Travis WD, Adusumilli PS. Prognostic immune markers in non-small cell lung cancer. *Clin Cancer Res.* 2011;17(16):5247-56.

Kadota K, Suzuki K, Sima CS, Rusch VW, Adusumilli PS, Travis WD. Pleomorphic epithelioid diffuse malignant pleural mesothelioma: a clinicopathological review and conceptual proposal to reclassify as biphasic or sarcomatoid mesothelioma. *J Thorac Oncol.* 2011;6(5):896-904.

▼ Connect

[Contact us](#)

[Locations](#)

APPOINTMENTS

[800-525-2225](#)



▼ About MSK

[About us](#)

[Careers](#) 

[Giving](#) 

▾ Cancer Care

[Adult cancer types](#)

[Child & teen cancer types](#)

[Integrative medicine](#)

[Nutrition & cancer](#)

[Find a doctor](#)

▾ Research & Education

[Sloan Kettering Institute](#)

[Gerstner Sloan Kettering Graduate School](#) 

[Graduate medical education](#)

[MSK Library](#) 

[Communication preferences](#)

[Cookie preferences](#)

[Legal disclaimer](#)

[Accessibility statement](#)

[Privacy policy](#)

[Price transparency](#)

[Public notices](#)

© 2024 Memorial Sloan Kettering Cancer Center