



PATIENT & CAREGIVER EDUCATION

Alitreinoin (Systemic

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: Canada

Toctino [DSC]

Warning

- Do not take if you are pregnant or if you may get pregnant. The risk of very bad and sometimes deadly birth defects is very high if you take this drug at any time while you are pregnant. Any unborn baby can be harmed. There is no good way to tell if an unborn baby has been harmed. The risk of losing an unborn baby is also raised, and premature births have happened. Your doctor will talk about the bad effects before starting you on this drug. If you get pregnant while taking this drug or within 1 month after you stop taking it, call your doctor right away. If you know all the facts

and can follow how to take this drug you must sign a patient fact/consent form. Do not sign the form and do not take this drug if you do not know everything on the form.

- You must have 2 pregnancy tests that show you are NOT pregnant before starting this drug. You must have pregnancy tests done while taking this drug and 1 month after the last dose. If you have questions, talk with your doctor.
- Use 2 kinds of birth control 1 month before starting this drug, during treatment, and for at least 1 month after the last dose.
- Raised pressure in the brain has happened with this drug. This can cause long lasting loss of eyesight and sometimes death. Call your doctor right away if you have a bad headache, dizziness, upset stomach or throwing up, or seizures. Call your doctor right away if you have weakness on 1 side of the body, trouble speaking or thinking, change in balance, or change in eyesight.

What is this drug used for?

- It is used to treat hand eczema.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you are allergic to soya lecithin or alike food products such as soybeans or peanuts, talk with the doctor.
- If you have any of these health problems: High cholesterol, high triglycerides, kidney disease, liver disease, or thyroid disease.
- If you have too much vitamin A in the blood.
- If you are not able to break down fructose.
- If your eczema can be treated by something else like other drugs.
- If you are taking any of these drugs: Demeclocycline, doxycycline, minocycline, tetracycline, a product that has vitamin A in it, a product that is like vitamin A, or St. John's wort.
- If you are taking any of these drugs: Amiodarone, isotretinoin, or tazarotene.
- If you are able to get pregnant and are not using 2 kinds of birth control.

- If you are planning to get pregnant within 1 month before treatment, during treatment, or within 1 month after your last dose.
- If you are breast-feeding. Do not breast-feed while you take this drug.
- If the patient is a child. Do not give this drug to a child.

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Avoid driving and doing other tasks or actions that call for you to be alert or have clear eyesight until you see how this drug affects you.

- Lowered night eyesight may happen. This may be sudden. This may clear up after you stop the drug but sometimes it may not go away.
- Use care when driving at night or doing other tasks in lowered lighting.
- It is common to have dry eyes with this drug. Artificial tears, lubricant eye gels, or other products may be used to prevent or treat dry eyes. If you have questions, talk with the doctor or pharmacist.
- Dry skin and lips are common with this drug. Skin moisturizers and lip balms may be used as the doctor has told you.
- If you have high blood sugar (diabetes), this drug may sometimes raise blood sugar. Talk with your doctor about how to keep your blood sugar under control.
- Do not donate blood while using this drug and for 1 month after stopping.
- Treatment with this drug may lead to higher cholesterol and triglycerides. The effect of these changes on heart health is not known. Talk with the doctor.
- Have your blood work and other lab tests checked as you have been told by your doctor.
- Talk with your doctor before you drink alcohol.

- Skin may look worse before it looks better.
- You may get sunburned more easily. Avoid sun, sunlamps, and tanning beds. Use sunscreen and wear clothing and eyewear that protects you from the sun.
- This drug may cause weak bones and tendon problems in some people. The chance of bone problems like broken bones may be raised in people who play certain sports. Talk with the doctor.
- This drug may affect being able to father a child. Talk with the doctor.
- Do not use progestin-only birth control pills (minipills). They may not work well. Talk with your doctor.
- If you have sex without using 2 kinds of birth control that you can trust, if you think you may be pregnant, or if you miss your period, call your doctor right away.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Signs of high blood sugar like confusion, feeling sleepy, unusual thirst or hunger, passing urine more often, flushing, fast breathing, or breath that smells like fruit.
- Signs of depression, suicidal thoughts, emotional ups and downs, abnormal thinking, anxiety, or lack of interest in life.
- Behavior problems.
- Back, bone, joint, or muscle pain.
- Change in eyesight, eye pain, or severe eye irritation.
- Change in how contact lenses feel in the eyes.
- Any unexplained bruising or bleeding.
- A pancreas problem (pancreatitis) has happened with

this drug. Pancreatitis can be deadly. Call your doctor right away if you have severe stomach pain, back pain, or upset stomach or throwing up.

- Severe bowel problems may happen with this drug (inflammatory bowel disease). Tell your doctor right away if you have signs like severe diarrhea, stomach pain, bleeding from the rectum, or rectal pain. This may clear up after you stop the drug but sometimes it may not go away.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Headache.
- Flushing.
- Redness.
- Dry mouth, eyes, skin, or lips.
- Nose or throat irritation.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

- Take with a meal.
- Swallow whole. Do not chew, break, or crush.
- Take this drug at the same time of day.
- Use as you have been told, even if your signs get better.

What do I do if I miss a dose?

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.
- Do not take 2 doses on the same day.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Store in the original container to protect from light.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your

doctor, nurse, pharmacist, or other health care provider.

- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate,

Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2023-02-27

Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Alitretinoin (Systemic - Last updated on December 12, 2022
All rights owned and reserved by Memorial Sloan Kettering
Cancer Center