

PATIENT & CAREGIVER EDUCATION

Diclofenac (Topical)

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Aspercreme Arthritis Pain [OTC]; Diclogen; Dicloprep-100 [OTC] [DSC]; Diclotrex II [DSC]; Diclotrex [DSC]; Flector; FT Arthritis Pain [OTC]; Licart; Lixofen; Motrin Arthritis Pain [OTC]; Pennsaid [DSC]; Pharmacist Choice Diclofenac [OTC]; Profinac; ValcoPrep-100 [DSC]; VennGel One; VennGel Two; Voltaren Arthritis Pain [OTC]; Xiclo; Xrylix II [DSC]; Xrylix [DSC]

Brand Names: Canada

JAMP Diclofenac; Pennsaid; PMS-Diclofenac; TARO-Diclofenac

Warning

- This drug may raise the chance of heart and blood vessel side effects like heart attack and stroke. If these happen, they can be deadly. The risk of these side effects may be greater if your child has heart disease or risks for heart disease. However, the risk may also be raised in people who do not have heart disease or risks for heart disease. The risk of these health problems can happen as soon as the first weeks of using this drug and may be greater with higher doses or with long-term use. Do not give this drug to your child right before or after bypass heart surgery.
- This drug may raise the chance of severe and sometimes deadly stomach or bowel problems like ulcers or bleeding. The risk is greater in older people, and in people who have had stomach or bowel ulcers or bleeding before. These problems may occur without warning signs.

What is this drug used for?

Skin patch:

- It is used to manage pain.
- It is used to treat some types of arthritis.

Cream, gel, or topical solution:

- This form of this drug is not approved for use in children. However, your child's doctor may decide the benefits of taking this drug outweigh the risks. If your child has been given this form of this drug, ask your child's doctor for information about the benefits and risks. Talk with your child's doctor if you have questions about this drug.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has an allergy to aspirin or nonsteroidal anti-inflammatory drugs (NSAIDs) like ibuprofen or naproxen.
- If your child has ever had asthma caused by a salicylate drug like aspirin or a drug like this one like NSAIDs.
- If your child has any of these health problems: Dehydration, GI (gastrointestinal) bleeding, heart failure (weak heart), kidney disease, or liver disease.

- If your child has had a recent heart attack.
- If your child is taking any other NSAID, a salicylate like aspirin, or pemetrexed.
- If your child is having their fertility checked.

If your child is pregnant:

- Talk to the doctor if your child is pregnant, plans to become pregnant, or gets pregnant while taking this drug. This drug may cause harm to an unborn baby if taken at 20 weeks or later in pregnancy. If your child is between 20 to 30 weeks of pregnancy, only give this drug if the doctor has told you to. Do not give this drug to your child if she is more than 30 weeks pregnant.

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

Skin patch:

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have blood work checked as you have been told by the doctor. Talk with the doctor.
- High blood pressure has happened with drugs like this one. Have your child's blood pressure checked as you have been told by the doctor.
- If your child smokes, talk with the doctor.
- If your child has asthma, talk with the doctor. Your child may be more sensitive to this drug.
- Alcohol may interact with this drug. Be sure your child does not drink alcohol.
- Do not use on skin that has any problems.
- Do not use more than told. Unsafe side effects may happen.
- Do not use longer than you have been told by the doctor.
- The chance of heart failure is raised with the use of drugs like this one. In people who already have heart failure,

the chance of heart attack, having to go to the hospital for heart failure, and death is raised. Talk with the doctor.

- The chance of heart attack and heart-related death is raised in people taking drugs like this one after a recent heart attack. People taking drugs like this one after a first heart attack were also more likely to die in the year after the heart attack compared with people not taking drugs like this one. Talk with the doctor.
- If your child is taking aspirin to help prevent a heart attack, talk with the doctor.
- This drug may cause harm if chewed or swallowed. If this drug has been put in the mouth, call a doctor or poison control center right away.

Skin patch 1.25%:

- Your child's skin may be sensitive to heat for a few days after using this drug. This includes from heat sources like hot showers or baths, sunlight, sunlamps, tanning beds, heating pads, electric blankets, heat lamps, saunas, hot tubs, and heated waterbeds. It also includes heat from exercise.
- Your child may need to avoid sun, sunlamps, and tanning beds on the treated area for some time after taking off the patch. Talk with the doctor.

If your child is or may be sexually active:

- NSAIDs like this drug may affect egg release (ovulation). This may affect being able to get pregnant. This goes back to normal when this drug is stopped. Talk with the doctor.

If your child is breast-feeding a baby:

- Tell the doctor if your child is breast-feeding a baby. You will need to talk about any risks to the baby.

What are some side effects that I need to call my child's doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of bleeding like throwing up or coughing up blood; vomit that looks like coffee grounds; blood in the urine;

black, red, or tarry stools; bleeding from the gums; abnormal vaginal bleeding; bruises without a cause or that get bigger; or bleeding you cannot stop.

- Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Signs of high potassium levels like a heartbeat that does not feel normal; feeling confused; feeling weak, lightheaded, or dizzy; feeling like passing out; numbness or tingling; or shortness of breath.
- Skin irritation.
- Chest pain or pressure or a fast heartbeat.
- Shortness of breath, a big weight gain, or swelling in the arms or legs.
- Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight.
- Feeling very tired or weak.
- Flu-like signs.
- Very bad back pain.
- Severe skin reactions may happen with this drug. These

include Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and other serious reactions. Sometimes, body organs may also be affected. These reactions can be deadly. Get medical help right away if your child has signs like red, swollen, blistered, or peeling skin; red or irritated eyes; sores in the mouth, throat, nose, eyes, genitals, or any areas of skin; fever; chills; body aches; shortness of breath; or swollen glands.

- Liver problems have happened with drugs like this one. Sometimes, this has been deadly. Call the doctor right away if your child has signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Dizziness or headache.
- Constipation, diarrhea, stomach pain, upset stomach, or throwing up.

- Gas.
- Heartburn.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

Skin patch:

- Do not give by mouth. Use on your child's skin only. Keep out of your child's mouth, nose, and eyes (may burn).
- Wash your hands before and after use.
- Put patch on clean, dry, healthy skin. Move the site with each new patch.
- Clean affected part before use. Make sure to dry well.
- Do not put on cuts, scrapes, eczema, or damaged skin.
- Do not put on skin where you have already used cosmetics, lotions, insect repellant, sunscreen, or other

skin products.

- Do not let your child bathe, shower, or swim after putting on.
- Have your child avoid use of heat sources (such as sunlamps, tanning beds, heating pads, electric blankets, heat lamps, saunas, hot tubs, heated waterbeds). Avoid long, hot baths or sunbathing. Your child's temperature may rise and cause too much drug to pass into your child's body.
- If the patch does not stick well, talk with your pharmacist about what to do. Certain things can be done to help hold it in place.
- If the patch falls off, put a new one on.
- If you get this drug in your child's eyes, wash right away with water. If your child has eye irritation that lasts or a change in eyesight, call the doctor.

Skin patch 1.3% (Flector, Licart):

- If the patch loosens, put tape ONLY on the edges of the patch to hold it in place.

Skin patch 1.25% (Xiclo):

- One patch may be used 2 times in the same day. If you take this patch off and want to keep it for later, place the

clear plastic film back onto the sticky part of the patch. If you have questions about how to reuse this patch, talk to the pharmacist. Be sure you store the used patch out of the reach of children and pets.

- Do not leave this drug on the skin longer than you have been told by the doctor.
- Your child may wear clothing over the area where this drug is placed.

What do I do if my child misses a dose?

- Skip the missed dose and go back to your child's normal time unless your child's doctor tells you to do something else.
- Do not put on 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature.
- Protect from heat.
- Store in the pouch this drug comes in. Keep the pouch sealed when it is not in use. The pouch is not child proof.
- After opening, be sure you know how long the product is good for and how to store it. Ask the doctor or pharmacist if you are not sure.
- After you take off a skin patch, be sure to fold the sticky

sides of the patch to each other. Throw away used patches where children and pets cannot get to them.

- Throw away unused patches when they are no longer needed. Take them from the pouch, take off liner, and fold the sticky side of the patch together.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child's doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready

to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical->

Last Reviewed Date

2025-03-24

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors.
All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Diclofenac (Topical - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering
Cancer Center