MSK Kids

PATIENT & CAREGIVER EDUCATION

Ixekizumab

This information from Lexicomp® explains what you need to know about this medication, including what it’s used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Taltz

Brand Names: Canada

Taltz

What is this drug used for?

- It is used to treat plaque psoriasis.
- It may be given to your child for other reasons. Talk with the doctor.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.
- If your child has TB (tuberculosis).

This is not a list of all drugs or health problems that interact with this drug.

Tell the doctor and pharmacist about all of your child’s drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child’s other drugs and health problems. Do not start, stop, or change the dose of any drug your

child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child’s health care providers that your child is taking this drug. This includes your child’s doctors, nurses, pharmacists, and dentists.
- Your child will need a TB (tuberculosis) test before starting this drug.
- Your child may have more chance of getting infections. Have your child wash hands often. Avoid crowds and people with infections, colds, or flu.
- If your child has inflammatory bowel disease like ulcerative colitis or Crohn’s disease, talk with the doctor. New and worsening inflammatory bowel disease has happened with this drug. Sometimes this may be severe. If you have questions, talk with the doctor.
- Make sure your child is up to date with all vaccines before treatment with this drug.
- Talk with the doctor before your child gets any vaccines. Use of some vaccines with this drug may either raise the chance of very bad infection or make the vaccine not work as well.
- Allergic reactions have happened with this drug. Sometimes, these have been very bad and people have had to go to the hospital. Talk with the doctor.
- The dose of this drug is based on your child’s weight. Some doses need to be given by a doctor or other health care provider. If you have questions, talk with the doctor.

If your child is pregnant or breast-feeding a baby:

- Talk with the doctor if your child is pregnant, becomes pregnant, or is breast-feeding a baby. You will need to talk about the benefits and risks to your child and the baby.

What are some side effects that I need to call my child’s doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child’s doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal.
- Warm, red, or painful skin or sores on the body.
- Redness or white patches in mouth or throat.
- Eye redness.
- Stomach pain or diarrhea.
- Bloody diarrhea.
- Weight loss.
- Very bad dizziness or passing out.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child’s doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Irritation where the shot is given.
- Upset stomach.
- Signs of a common cold.

These are not all of the side effects that may occur. If you have questions about side effects, call your child’s doctor. Call your child’s doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child’s doctor. Read all information given to you. Follow all instructions closely.

All products:

- It is given as a shot into the fatty part of the skin on the top of the thigh, belly area, or upper arm.
- If you will be giving your child the shot, your child’s doctor or nurse will teach you how to give the shot.
- Before using this drug, take it out of the refrigerator and leave it at room temperature for 30 minutes.
- Do not heat, microwave, place under hot water, or leave in direct sunlight.
- Wash your hands before and after use.
- Move the site where you give the shot with each shot.
- Do not shake.
- Do not give into skin that is irritated, bruised, red, infected, hard, or scarred.
- Do not give into skin that is affected by psoriasis.
- Do not give into skin within 1 inch of the belly button.
- Do not use if the solution is cloudy, leaking, or has particles.
- This drug is colorless to a faint yellow. Do not use if the solution changes color.
- Throw away any part left over after the dose is given.

- Throw away needles in a needle/sharp disposal box. Do not reuse needles or other items. When the box is full, follow all local rules for getting rid of it. Talk with a doctor or pharmacist if you have any questions.

Auto-injector shot:

- If you have trouble hearing or seeing, have someone else give the shot.

Prefilled syringes:

- If you have trouble seeing, have someone else give the shot.

What do I do if my child misses a dose?

- Give a missed dose as soon as you think about it and go back to your child’s normal time.
- If it is close to the time for your child’s next dose, skip the missed dose and go back to your child’s normal time.
- Do not give 2 doses at the same time or extra doses.
- If you are not sure what to do if your child misses a dose, call the doctor.

How do I store and/or throw out this drug?

All products:

- Store in a refrigerator. Do not freeze.
- Do not use if it has been frozen.
- Store in the original container to protect from light.
- If needed, you may store this drug in the original carton at room temperature for up to 5 days. Write down the date you take this drug out of the refrigerator. If stored at room temperature and not used within 5 days, throw this drug away.
- Do not put this drug back in the refrigerator after it has been stored at room temperature.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children

and pets.

- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

Auto-injector shot:

- If this drug comes in a pack with more than 1 auto-injector, take 1 auto-injector out of the original carton at a time. Protect it from light. Leave the other auto-injectors in the original carton in the refrigerator.
- If you drop this drug on a hard surface, do not use it.

General drug facts

- If your child’s symptoms or health problems do not get better or if they become worse, call your child’s doctor.
- Do not share your child’s drug with others and do not give anyone else’s drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child’s doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based

on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2022-10-17

Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Ixekizumab - Last updated on December 12, 2022
All rights owned and reserved by Memorial Sloan Kettering Cancer Center

Memorial Sloan Kettering Cancer Center