MSK Kids

PATIENT & CAREGIVER EDUCATION

Tiopronin

This information from Lexicomp® explains what you need to know about this medication, including what it’s used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Thiola; Thiola EC

What is this drug used for?

- It is used to prevent kidney stones.

What do I need to tell the doctor BEFORE my child takes this drug?

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.

If your child is breast-feeding a baby:

- Be sure your child does not breast-feed a baby while taking this drug.

This drug may interact with other drugs or health problems.

Tell the doctor and pharmacist about all of your child’s drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child’s other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

What are some things I need to know or do while my child takes this drug?

- Tell all of your child’s health care providers that your child is taking this drug. This includes your child’s doctors, nurses, pharmacists, and dentists.
- Have your child’s blood work and other lab tests checked as you have been told by the doctor.
- Alcohol may interact with this drug. Be sure your child does not drink alcohol.

If your child is pregnant:

- Tell the doctor if your child is pregnant or becomes pregnant. You will need to talk about the benefits and risks of your child using this drug while pregnant.

What are some side effects that I need to call my child’s doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child’s doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Fever.
- Joint pain.
- Swollen gland.

- Any skin change.
- Swelling in the arms or legs.
- Chest pain.

If your child is or may be sexually active:

- Not able to get or keep an erection.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child’s doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Diarrhea, stomach pain, upset stomach, throwing up, or decreased appetite.
- Change in taste.
- Mouth sores.
- Cough.
- Itching.
- Feeling tired or weak.

These are not all of the side effects that may occur. If you have questions about side effects, call your child’s doctor. Call your child’s doctor for medical advice about side effects.

You may report side effects to your national health agency.

How is this drug best given?

Give this drug as ordered by your child’s doctor. Read all information given to you. Follow all instructions closely.

Regular-release tablets:

- Give on an empty stomach at least 1 hour before or 2 hours after meals unless the doctor has told you otherwise.

Delayed-release tablets:

- Give this drug with or without food but give it the same way each time. Always give with food or always give on an empty stomach.
- If your child cannot swallow tablets whole, you may crush the tablets and mix with applesauce. Crush 1 tablet at a time. If needed, the mixture may be stored in the refrigerator for 2 hours after mixing. After giving the dose, add water to the container, mix, and have your child drink.

All products:

- Give this drug at the same time of day.
- Have your child follow the diet plan your child’s doctor told you about.
- Follow what your child’s doctor has told you about the type of liquids and how much liquid your child can drink while taking this drug.

What do I do if my child misses a dose?

- Give a missed dose as soon as you think about it.
- If it is close to the time for your child’s next dose, skip the missed dose and go back to your child’s normal time.
- Do not give 2 doses at the same time or extra doses.

How do I store and/or throw out this drug?

- Store at room temperature in a dry place. Do not store in a bathroom.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if

you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your child’s symptoms or health problems do not get better or if they become worse, call your child’s doctor.
- Do not share your child’s drug with others and do not give anyone else’s drug to your child.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your child’s doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider’s examination and assessment of a patient’s specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms.

Last Reviewed Date

2024-01-24

Copyright

© 2024 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have any questions, contact a member of your care team directly. If you're a patient at MSK and you need to reach a provider after 5 p.m., during the weekend, or on a holiday, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Tiopronin - Last updated on December 12, 2022
All rights owned and reserved by Memorial Sloan Kettering Cancer Center

Memorial Sloan Kettering Cancer Center