

## Tisagenlecleucel

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

### Brand Names: US

Kymriah

### Brand Names: Canada

Kymriah

## Warning

- A severe health problem called cytokine release syndrome (CRS) has happened with this drug. Sometimes, this has been life-threatening or deadly. Get medical help right away if your child has signs like chills; confusion; dizziness; fast or abnormal heartbeat; chest pain; change in how much urine is passed; feeling tired or weak; fever; headache; passing out; rash; swelling; trouble breathing; severe muscle or joint pain; yellow skin or eyes; dark urine; severe upset stomach, throwing up, or diarrhea; or wheezing.
- If your child has an infection or any inflammatory health problem, talk with your child's doctor. This drug may not be right for your child.
- Nervous system problems have happened after treatment with this drug. Sometimes, these have been severe and life-threatening or deadly. Call your child's doctor right away if your child feels agitated, confused, nervous, sluggish, or restless. Call your child's doctor right away if your child has anxiety, change in balance, change in eyesight, dizziness or passing out, hallucinations (seeing or hearing things that are not there), headaches, seizures, shakiness, trouble speaking, trouble reading or writing, or new or worse behavior or mood changes.
- You will need to stay close to where your child got this drug for your child to be watched for signs of CRS and nervous system problems for at least 2 weeks after getting it.
- This drug may raise the risk of other cancers. This includes certain types of blood cancer. The doctor will watch for this. If you have questions, talk with the doctor.

## What is this drug used for?

- It is used to treat a type of leukemia.
- If your child has been given this drug for some other reason, talk with the doctor about the benefits and risks. Talk with the doctor if you have

questions or concerns about giving this drug to your child.

## **What do I need to tell the doctor BEFORE my child takes this drug?**

- If your child is allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell the doctor about the allergy and what signs your child had.

### **If your child is pregnant:**

- Do not give this drug to your child during pregnancy.

This drug may interact with other drugs or health problems.

Tell the doctor and pharmacist about all of your child's drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe to give this drug with all of your child's other drugs and health problems. Do not start, stop, or change the dose of any drug your child takes without checking with the doctor.

## **What are some things I need to know or do while my child takes this drug?**

- Tell all of your child's health care providers that your child is taking this drug. This includes your child's doctors, nurses, pharmacists, and dentists.
- Have your child avoid tasks or actions that call for alertness for at least 2 weeks after getting this drug. If you have any questions, call your child's doctor or pharmacist.
- Have your child's blood work and other lab tests checked as you have been told by the doctor.
- This drug may affect certain lab tests. Tell all of your child's health care providers and lab workers that your child takes this drug.
- Low blood cell counts have happened with this drug. This may last for several weeks after taking this drug. If blood cell counts get very low, this

can lead to bleeding problems, infections, or anemia. Some severe infections have been deadly. If you have questions, talk with the doctor.

- Have your child wash hands often. Avoid crowds and people with infections, colds, or flu.
- Your child may bleed more easily. Make sure your child is careful and avoids injury. Be sure your child has a soft toothbrush.
- If your child has had hepatitis B before or carries the virus, talk with your child's doctor. Drugs like this one can cause the virus to become active. This can lead to very bad and sometimes deadly liver problems.
- Hepatitis and HIV testing will be done before taking this drug.
- Talk with the doctor before your child gets any vaccines. Some types of vaccines must not be given for at least 6 weeks before starting this drug, during treatment, and for some time after the last dose.
- Do not let your child donate blood, organs, or tissues or cells. Talk with your child's doctor to see when your child can do these things after getting this drug.
- If your child has an upset stomach or diarrhea, is throwing up, or decreased appetite, talk with the doctor. There may be ways to lower these side effects.
- If your child is of childbearing age, a pregnancy test will need to be done before starting this drug to make sure your child is not pregnant.

### **If your child is or may be sexually active:**

- This drug may cause harm to the unborn baby if your child takes it during pregnancy.
- Have your child use birth control while taking this drug and for some time after the last dose. Ask the doctor how long your child must use birth control. If your child becomes pregnant, call the doctor right away.
- If your child wants to get pregnant after stopping this drug, talk with the doctor.

## If your child is breast-feeding a baby:

- Tell the doctor if your child is breast-feeding a baby. You will need to talk about any risks to the baby.

## What are some side effects that I need to call my child's doctor about right away?

**WARNING/CAUTION:** Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your child's doctor or get medical help right away if your child has any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal.
- Signs of bleeding like throwing up or coughing up blood; vomit that looks like coffee grounds; blood in the urine; black, red, or tarry stools; bleeding from the gums; abnormal vaginal bleeding; bruises without a cause or that get bigger; or bleeding you cannot stop.
- Signs of liver problems like dark urine, tiredness, decreased appetite, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.
- Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain.
- Signs of high or low blood pressure like very bad headache or dizziness, passing out, or change in eyesight.
- Signs of electrolyte problems like mood changes; confusion; muscle pain, cramps, or spasms; weakness; shakiness; change in balance; an abnormal heartbeat; seizures; loss of appetite; or severe upset stomach or throwing

up.

- Signs of a severe health problem called tumor lysis syndrome (TLS) like fast or abnormal heartbeat; passing out; trouble passing urine; muscle weakness or cramps; upset stomach, throwing up, diarrhea, or not able to eat; or feeling sluggish.
- Shortness of breath, a big weight gain, or swelling in the arms or legs.
- Fast breathing.
- Fast or abnormal heartbeat.
- Trouble sleeping.
- Feeling very sleepy.
- Not able to focus.
- Memory problems or loss.
- Swelling.
- Trouble moving around.
- Bad dreams.
- A burning, numbness, or tingling feeling that is not normal.
- Call the doctor right away if your child has signs of a blood clot like chest pain or pressure; coughing up blood; shortness of breath; swelling, warmth, numbness, change of color, or pain in a leg or arm; or trouble speaking or swallowing.
- An immune system problem called hemophagocytic lymphohistiocytosis/macrophage activation syndrome (HLH/MAS) has happened with this drug. HLH/MAS can be life-threatening or deadly. Call the doctor right away if your child feels confused or not alert or has fever, swollen gland, rash, seizures, change in balance, or trouble walking that is new or worse.
- A severe brain problem called progressive multifocal leukoencephalopathy (PML) has happened with this drug. It may cause disability or can be deadly. Tell the doctor right away if your child has signs like confusion, memory problems, depression, change in the way your child acts, change in strength

on 1 side is greater than the other, trouble speaking or thinking, change in balance, or change in eyesight.

## What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your child's doctor or get medical help if any of these side effects or any other side effects bother your child or do not go away:

- Constipation, diarrhea, stomach pain, upset stomach, throwing up, or decreased appetite.
- Headache.
- Feeling tired or weak.
- Pain in arms or legs.
- Back, bone, joint, muscle, or neck pain.
- Stuffy nose.
- Weight loss.

These are not all of the side effects that may occur. If you have questions about side effects, call your child's doctor. Call your child's doctor for medical advice about side effects.

You may report side effects to your national health agency.

## How is this drug best given?

Give this drug as ordered by your child's doctor. Read all information given to you. Follow all instructions closely.

- It is given as an infusion into a vein over a period of time.
- Other drugs may be given before this drug to help avoid side effects.

## What do I do if my child misses a dose?

- Call your child's doctor to find out what to do.

## How do I store and/or throw out this drug?

- This drug will be given in a hospital or doctor's office. You will not store it at home.

## General drug facts

- If your child's symptoms or health problems do not get better or if they become worse, call your child's doctor.
- Do not share your child's drug with others and do not give anyone else's drug to your child.
- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

## Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific

patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at <https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

## Last Reviewed Date

2026-01-27

## Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit [www.mskcc.org/pe](https://www.mskcc.org/pe) to search our virtual library.

Tisagenlecleucel - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center



Memorial Sloan Kettering  
Cancer Center