



PATIENT & CAREGIVER EDUCATION

Propranolol

This information from Lexicomp® explains what you need to know about this medication, including what it's used for, how to take it, its side effects, and when to call your healthcare provider.

Brand Names: US

Hemangeol; Inderal LA; Inderal XL; InnoPran XL

Brand Names: Canada

APO-Propranolol [DSC]; Hemangiol; Inderal LA [DSC]; LUPIN-Propranolol LA; PRZ-Propranolol; TEVA-Propranolol

What is this drug used for?

- It is used to treat high blood pressure.
- It is used to treat chest pain or pressure.
- It is used to help certain heart problems.
- It is used to prevent migraine headaches.
- It is used to treat tremor (essential).
- It is used after a heart attack to help prevent future heart attacks and lengthen life.
- It is used to treat pheochromocytoma.
- It is used to treat certain types of abnormal heartbeats.
- It may be given to you for other reasons. Talk with the doctor.

What do I need to tell my doctor BEFORE I take this drug?

- If you are allergic to this drug; any part of this drug; or any other drugs, foods, or substances. Tell your doctor about the allergy and what signs you had.
- If you have any of these health problems: Certain types of abnormal heartbeats called heart block or sick-sinus syndrome, heart failure (weak heart), low blood pressure, poor blood flow to the arms or legs, shock caused by heart problems, or a slow heartbeat.
- If you have any of these health problems: Asthma or other breathing problems like COPD (chronic obstructive pulmonary disease).

This is not a list of all drugs or health problems that interact with this drug.

Tell your doctor and pharmacist about all of your drugs (prescription or OTC, natural products, vitamins) and health problems. You must check to make sure that it is safe for you to take this drug with all of your drugs and health problems. Do not start, stop, or change the dose of any drug without checking with your doctor.

What are some things I need to know or do while I take this drug?

All products:

- Tell all of your health care providers that you take this drug. This includes your doctors, nurses, pharmacists, and dentists.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this drug affects you.
- To lower the chance of feeling dizzy or passing out, rise slowly if you have been sitting or lying down. Be careful going up and down stairs.
- Check blood pressure and heart rate as the doctor has told you.
- This drug may affect certain lab tests. Tell all of your health care providers and lab workers that you take this drug.

- Talk with your doctor before you drink alcohol.
- If you smoke, talk with your doctor.
- This drug may prevent some signs of low blood sugar like fast heartbeat. This may raise the risk of severe or long-lasting low blood sugar, especially in people with diabetes, children, and people who are fasting. This includes people who are having surgery, are not eating like normal, or are throwing up. If you have questions, talk with the doctor.
- If you have high blood sugar (diabetes), you will need to watch your blood sugar closely.
- Do not stop taking this drug all of a sudden. If you do, chest pain that is worse and in some cases heart attack may occur. The risk may be greater if you have certain types of heart disease. To avoid side effects, you will want to slowly stop this drug as ordered by your doctor. Call your doctor right away if you have new or worse chest pain or if other heart problems occur.
- This drug may make it harder to tell if you have signs of an overactive thyroid like fast heartbeat. If you have an overactive thyroid and stop taking this drug all of a sudden, it may get worse and could be life-threatening. Talk with your doctor.
- If you are taking this drug and have high blood pressure, talk with your doctor before using OTC products that may raise blood pressure. These include cough or cold drugs, diet pills, stimulants, non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen or naproxen, and some natural products or aids.
- If you have had a very bad allergic reaction, talk with your doctor. You may have a chance of an even worse reaction if you come into contact with what caused your allergy. If you use epinephrine to treat very bad allergic reactions, talk with your doctor. Epinephrine may not work as well while you are taking this drug.
- Tell your doctor if you are pregnant, plan on getting pregnant, or are breast-feeding. You will need to talk about the benefits and risks to you and the baby.

Oral solution:

- Make sure you have the right liquid; there is more than one strength.

What are some side effects that I need to call my doctor about right away?

WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect:

- Signs of an allergic reaction, like rash; hives; itching; red, swollen, blistered, or peeling skin with or without fever; wheezing; tightness in the chest or throat; trouble breathing, swallowing, or talking; unusual hoarseness; or swelling of the mouth, face, lips, tongue, or throat.
- Signs of low blood sugar like dizziness, headache, feeling sleepy, feeling weak, shaking, a fast heartbeat, confusion, hunger, or sweating.
- Signs of lupus like a rash on the cheeks or other body parts, sunburn easy, muscle or joint pain, chest pain or shortness of breath, or swelling in the arms or legs.
- Very bad dizziness or passing out.
- Chest pain that is new or worse.
- Slow heartbeat.
- A heartbeat that does not feel normal.
- Shortness of breath, a big weight gain, or swelling in the arms or legs.
- Feeling confused.
- Hallucinations (seeing or hearing things that are not there).
- Memory problems or loss.
- Depression or other mood changes.
- A burning, numbness, or tingling feeling that is not normal.
- Feeling cold in the arms or legs.

- Change in color of hands, feet, or other areas. Skin may turn pale, blue, gray, purple, or red.
- Change in eyesight.
- Any unexplained bruising or bleeding.
- Fever, chills, or sore throat.
- Not able to get or keep an erection.
- Severe skin reactions have happened with this drug. These have included Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and other severe skin reactions. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin; other skin irritation (with or without fever); red or irritated eyes; or sores in your mouth, throat, nose, or eyes.

What are some other side effects of this drug?

All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away:

- Feeling dizzy, sleepy, tired, or weak.
- Upset stomach or throwing up.
- Stomach pain or cramps.
- Diarrhea or constipation.
- Trouble sleeping.
- Strange or odd dreams.

These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects.

You may report side effects to your national health agency.

You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>.

How is this drug best taken?

Use this drug as ordered by your doctor. Read all information given to you. Follow all instructions closely.

Inderal XL and InnoPran XL:

- Take with or without food but take the same way each time. Always take with food or always take on an empty stomach.
- Take at bedtime if taking once a day.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.

All other oral products:

- Some drugs may need to be taken with food or on an empty stomach. For some drugs it does not matter. Check with your pharmacist about how to take this drug.
- Keep taking this drug as you have been told by your doctor or other health care provider, even if you feel well.

Oral solution:

- Measure liquid doses carefully. Use the measuring device that comes with this drug. If there is none, ask the pharmacist for a device to measure this drug.

All long-acting products:

- Swallow capsule whole. Do not chew, break, or crush.

Injection:

- It is given as a shot into a vein.

What do I do if I miss a dose?

All oral products:

- Take a missed dose as soon as you think about it.
- If it is close to the time for your next dose, skip the missed dose and go back to your normal time.
- Do not take 2 doses at the same time or extra doses.

Injection:

- Call your doctor to find out what to do.

How do I store and/or throw out this drug?

All oral products:

- Store at room temperature. Do not freeze.
- Protect from heat, cold, and light.
- Keep lid tightly closed.
- Store in a dry place. Do not store in a bathroom.

Injection:

- If you need to store this drug at home, talk with your doctor, nurse, or pharmacist about how to store it.

All products:

- Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets.
- Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area.

General drug facts

- If your symptoms or health problems do not get better or if they become worse, call your doctor.
- Do not share your drugs with others and do not take anyone else's drugs.
- Some drugs may have another patient information leaflet. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- Some drugs may have another patient information leaflet. Check with your pharmacist. If you have any questions about this drug, please talk with your doctor, nurse, pharmacist, or other health care provider.
- If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened.

Consumer Information Use and Disclaimer

This generalized information is a limited summary of diagnosis, treatment, and/or medication information. It is not meant to be comprehensive and should be used as a tool to help the user understand and/or assess potential diagnostic and treatment options. It does NOT include all information about conditions, treatments, medications, side effects, or risks that may apply to a specific patient. It is not intended to be medical advice or a substitute for the medical advice, diagnosis, or treatment of a health care provider based on the health care provider's examination and assessment of a patient's specific and unique circumstances. Patients must speak with a health care provider for complete information about their health, medical questions, and treatment options, including any risks or benefits regarding use of medications. This information does not endorse any treatments or medications as safe, effective, or approved for treating a specific patient. UpToDate, Inc. and its affiliates disclaim any warranty or liability relating to this information or the use thereof. The use of this information is governed by the Terms of Use, available at

<https://www.wolterskluwer.com/en/know/clinical-effectiveness-terms>.

Last Reviewed Date

2024-01-03

Copyright

© 2026 UpToDate, Inc. and its affiliates and/or licensors. All rights reserved.

If you have questions or concerns, contact your healthcare provider. A member of your care team will answer Monday through Friday from 9 a.m. to 5 p.m. Outside those hours, you can leave a message or talk with another MSK provider. There is always a doctor or nurse on call. If you're not sure how to reach your healthcare provider, call 212-639-2000.

For more resources, visit www.mskcc.org/pe to search our virtual library.

Propranolol - Last updated on December 12, 2022

All rights owned and reserved by Memorial Sloan Kettering Cancer Center